



Frontotemporal dementia, music perception and social cognition share neurobiological circuits: A meta-analysis

Jochum J. van't Hooft^{a,*}, Yolande A.L. Pijnenburg^a, Sietske A.M. Sikkes^a, Philip Scheltens^a, Jacoba M. Spikman^b, Artur C. Jaschke^{c,d}, Jason D. Warren^e, Betty M. Tijms^a

^a Alzheimer Centre Amsterdam, Department of Neurology, Amsterdam UMC, Amsterdam, the Netherlands

^b Department of Clinical Neuropsychology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

^c Clinical Neuropsychology, VU University Amsterdam, Amsterdam, the Netherlands

^d Music Therapy, ArteZ University of the Arts, Enschede, the Netherlands

^e Dementia Research Centre, UCL Queen Square Institute of Neurology, University College London, United Kingdom

ARTICLE INFO

Keywords:

Frontotemporal dementia
Music perception
Social cognition
Ventral language pathway
Insula

ABSTRACT

Frontotemporal dementia (FTD) is a neurodegenerative disease that presents with profound changes in social cognition. Music might be a sensitive probe for social cognition abilities, but underlying neurobiological substrates are unclear. We performed a meta-analysis of voxel-based morphometry studies in FTD patients and functional MRI studies for music perception and social cognition tasks in cognitively normal controls to identify robust patterns of atrophy (FTD) or activation (music perception or social cognition). Conjunction analyses were performed to identify overlapping brain regions. In total 303 articles were included: 53 for FTD (n = 1153 patients, 42.5% female; 1337 controls, 53.8% female), 28 for music perception (n = 540, 51.8% female) and 222 for social cognition in controls (n = 5664, 50.2% female). We observed considerable overlap in atrophy patterns associated with FTD, and functional activation associated with music perception and social cognition, mostly encompassing the ventral language network. We further observed overlap across all three modalities in mesolimbic, basal forebrain and striatal regions. The results of our meta-analysis suggest that music perception and social cognition share neurobiological circuits that are affected in FTD. This supports the idea that music might be a sensitive probe for social cognition abilities with implications for diagnosis and monitoring.

1. Introduction

Frontotemporal dementia (FTD) is a neurodegenerative disease characterized by frontal and temporal lobar degeneration. It is the second most prevalent cause of early-onset dementia following Alzheimer's disease (Harvey et al. 2003; Ratnavalli et al. 2002). Contrary to other types of dementia, personality change and behavioral abnormalities are characteristics of FTD, which can occur even in the absence of cognitive decline. Since FTD is highly heterogeneous, both clinically and pathologically, it often poses a diagnostic challenge (Rascovsky et al. 2011). An early hallmark of FTD is a loss of appropriate interpersonal conduct due to a decline in social cognition abilities (Adenzato et al. 2010; Piguet et al. 2011). Social cognition is a multi-componential term used to describe cognitive processes related to the perception, understanding, and implementation of social cues (Van Overwalle et al. 2015; Suchy and Holdnack 2013) and comprises abilities such as emotion

recognition, mentalizing (or Theory of Mind) and empathy. Impairment of social cognition is not specific for FTD (Agustus et al. 2015; Bora et al. 2016; Gossink et al. 2018), as it is also a key symptom of psychiatric disorders such as autism and schizophrenia (Baron-Cohen et al. 2000; Corcoran et al. 1995; Langdon et al. 2002), and several acute neurological disorders (Buunk et al. 2017; May et al. 2017; Nijssse et al. 2019; Xiao et al. 2017). Despite the fact that social cognitive impairments are core symptoms in FTD (Rascovsky et al. 2011), and result in increased caregiver burden (Guevara et al. 2015; Hsieh et al. 2013), as of yet there is no gold standard for measuring social cognition abilities in the context of FTD diagnostics. On the other hand, FTD is a neuroanatomically well-defined disorder and as such may serve as a neurobiological model to understand the complex system involved in social cognition. Improved understanding of the neurobiological basis involved in social cognition and its impairment in FTD might aid in developing diagnostic tools across illness and disorders.

* Corresponding author at: Alzheimer Centre Amsterdam, Department of Neurology, Amsterdam UMC, PO Box 7057, 1007 MB Amsterdam, the Netherlands.
E-mail address: j.vanthoof@amsterdamumc.nl (J.J. van't Hooft).

Music is suggested to be a sensitive probe for social cognition abilities (Clark et al. 2014, 2015; Downey et al. 2013; P. D. Fletcher et al. 2013); various studies have found that both emotional music processing and social cognition are disturbed in FTD. For example, Downey et al. found that FTD patients have a specific inability to attribute mental states to music ('musical mentalising'), which correlated with tested performance on social cognition (TASIT) and reported empathy (Downey et al. 2013). Furthermore, brain regions involved in music emotion-recognition were found to be involved in Theory of Mind in FTD, which suggests a shared neurobiological circuit (Omar et al. 2011). Case-reports and explorative imaging studies also suggest that music perception, just like social cognition, is altered in patients with FTD, with patients and caregivers reporting changes in music perception, varying from musicophilia (enhanced appreciation of music) (Fletcher et al. 2013) to change of musical taste (Boeve and Geda 2001; Fletcher et al. 2013; Geroldi et al. 2000; Sacks 2007) and sound-aversion (Hardy et al. 2016). One study found worse performance on social cognition abilities in FTD patients with musicophilia (Fletcher et al. 2013). As such it could be hypothesized that music perception shares neurobiological correlates with social cognition and brain areas affected in FTD. So far, research has mostly investigated atrophy in FTD and functional activity in social cognition and music perception separately. In FTD a network is affected which includes frontoinsular, cingulate and striatal structures (Seeley et al. 2009; Zhou et al. 2010); in music perception the temporal cortices and various frontal regions were identified (Janata 2015; Limb 2006); and in social cognition various frontal, temporal and parietal brain regions are identified (Molenberghs et al. 2016; Schurz et al. 2014). Studies that investigated neural substrates for social cognition in FTD patients demonstrated associations with fronto-insulo-temporal atrophy (Couto et al. 2013; Eslinger et al. 2011). We hypothesized that these circuits of atrophy in FTD are also involved in music perception. More specifically, that these circuits would be most affected in FTD subtypes in whom behavioral symptoms would be most prominent. Furthermore, we hypothesized that social cognition subcomponents empathy and Theory of Mind would involve specific brain regions within this circuit.

In this meta-analysis we investigated whether brain areas showing atrophy in FTD show anatomical overlap with brain areas that are normally activated during music perception and social cognition tasks to investigate shared neurobiological circuits. Furthermore, we performed subgroup analyses in FTD and social cognition to specify for FTD subtypes and social cognition subcomponents.

2. Methods and materials

2.1. Literature search and screening strategy

The PRISMA guideline was used to perform this meta-analysis (Moher et al. 2009). The literature search was performed in April 2020. A search and selection of papers was conducted in the BrainMap database using Sleuth 3.0.3 software (Fox et al. 2005; Fox and Lancaster 2002; Laird et al. 2005) and in PubMed and Google Scholar. BrainMap is a database where MRI scan results are stored of both functional and structural neuroimaging studies (Fox and Lancaster 2002). Articles and experiments selection were performed with Sleuth 3.0.3, which is the search engine of BrainMap. Additional imaging coordinates of results from the PubMed and Google Scholar searches were manually added for inclusion in this meta-analysis. Separate literature searches were performed on the topics 'frontotemporal dementia', 'social cognition' and 'music perception'. Below we discuss the search and screening process of these topics in more detail for each modality. Papers were included when (a) structural studies compared patients with frontotemporal dementia with cognitively normal controls; (b) functional studies included subjects with normal vision, audition for music perception and without psychiatric or neurological diseases for music perception and social cognition (c) whole-brain unbiased voxelwise analysis was performed;

(e) stereotaxic coordinates in Talairach or Montreal Neurological Institute (MNI) coordinates were reported; (f) articles were written in English. For a full overview of the selection process see the flowcharts in the supplemental material.

2.2. Frontotemporal dementia

To determine which brain areas show atrophy in frontotemporal dementia a search was conducted in BrainMap's Voxel-Based Morphometry database. We selected articles archived under the paradigm frontotemporal dementia (Subject > Diagnosis > is > Frontotemporal Dementia: 15 papers), frontotemporal lobar degeneration (Subject > Diagnosis > is > Frontotemporal Lobar Degeneration: 12 papers), semantic dementia (Subject > Diagnosis > is > Semantic Dementia: 8 papers) and progressive nonfluent aphasia (Subject > Diagnosis > is > Progressive Nonfluent Aphasia: 9 papers). All FTD subtypes were included. This resulted in 14 articles for inclusion. A manual search in PubMed and Google Scholar was performed using the terms 'Voxel-based morphometry'; 'VBM'; 'semantic dementia'; 'primary progressive nonfluent aphasia'; 'frontotemporal dementia'; 'frontotemporal lobar degeneration'; 'FTD' and 'FTLD' and resulted in 44 additional articles for inclusion. Papers from the same patient cohort were excluded.

2.3. Music perception

To determine which brain areas are activated during music perception tasks a search was conducted in the functional BrainMap database. We selected articles archived under the experimental paradigm music comprehension (Experiments > Paradigm Class > is > Music Comprehension: 65 papers) and passive listening (Experiments > Paradigm Class > is > Passive Listening: 113 papers). This resulted in 9 articles for inclusion. A manual search on PubMed and Google Scholar was performed using the terms 'magnetic resonance imaging'; 'MRI'; 'fMRI'; 'music perception'; 'music comprehension' and 'music processing' and resulted in 19 additional papers for inclusion.

2.4. Social cognition

To determine which brain areas are activated during social cognition tasks a search was conducted in the functional BrainMap database. We selected articles archived under the experimental paradigm 'Theory of Mind' (Experiments > Paradigm Class > is > Theory of Mind: 86 papers), 'Social Cognition' (Experiments > Behavioral domain > is > Cognition > Social Cognition: 127 papers) and body language perception (Experiments > Paradigm Class > is > Emotional Body Language Perception: 8 papers). This resulted in 17 articles for inclusion. A manual search in PubMed and Google Scholar was performed using the terms 'magnetic resonance imaging'; 'MRI'; 'fMRI'; 'social cognition'; 'theory of mind'; 'perspective taking'; 'mentalizing'; 'mentalising'; 'empathy'; 'emotion recognition' and 'emotion perception' and resulted in 205 additional articles for inclusion.

2.5. Statistical analyses

The meta-analysis was carried out in BrainMap's GingerALE 3.0.2 (<http://brainmap.org/ale/index.html>). The procedure was as follows: (1) All coordinates were converted to Talairach space using the Lancaster transformation before being entered into the analysis (Lancaster et al. 2007). (2) Activation Likelihood Estimate (ALE) meta-analysis calculations were performed (Simon B. Eickhoff et al. 2009; Simon B Eickhoff et al. 2012; Turkeltaub et al. 2012). Activation foci in a given experiment are combined for each voxel, resulting in a modeled activation map that describes the convergence of results of sets of voxels in the brain. (3) Thresholding of the ALE scores based on the null-hypothesis (Simon B Eickhoff et al. 2012).

First, we examined modality specific effects of patterns of atrophy

involved in FTD and functional activation in cognitive normal subjects (music perception and social cognition). We performed three pairwise conjunction analyses to study the overlap between: 1. FTD and music perception; 2. FTD and social cognition; 3. Music perception and social cognition. Finally, we performed subgroup analyses on FTD subtypes (bvFTD, semantic dementia, progressive nonfluent aphasia) to study whether results were specific to any subtype, and we repeated analyses on social cognition including only studies on empathy and Theory of Mind to investigate how these studies influence the results. Statistical maps were thresholded using cluster-level family-wise error (FWE) correction $P < 0.05$ (S.B. Eickhoff et al. 2012).

3. Results

Our search terms resulted in a total of 3290 articles of which 369 for FTD, 685 for music perception and 2236 for social cognition. Fifty-three articles met the inclusion criteria for FTD, 28 for music perception and 222 for social cognition and were subjected to meta-analysis (Flowcharts 1, 2, 3 in the supplementary material), including a total of 8694 subjects: 1153 with FTD, 1337 controls (881 foci; Table 1; Table S1), 540 cognitively normal subjects for music perception (539 foci; Table 2; Table S2) and 5664 cognitively normal subjects for social cognition (3421 foci; Table 3; Table S3). Participants with FTD showed a slightly higher proportion of males whereas the controls were evenly distributed (42.5% and 53.8% female). Gender was evenly distributed amongst participants in music perception (51.8% female) and social cognition studies (50.2% female).

3.1. Modality specific meta-analytic results

3.1.1. Frontotemporal dementia atrophy correlates

The meta-analysis showed atrophy in FTD involving frontal structures such as the inferior, middle and superior frontal gyri, anterior cingulate gyri, temporal structures such as the insula, inferior, middle and superior temporal gyri and subcortical regions such as the amygdala and striatum (Fig. 1; Table S4). Repeating analyses separately for FTD subtypes, we observed specific bilateral frontoinsular and striatal atrophy patterns in bvFTD and semantic dementia and left-sided atrophy in PNFA. Additionally, bvFTD and semantic dementia showed atrophy in the amygdala. Semantic dementia and PNFA showed atrophy of the superior and middle temporal gyri (Fig. S1-3; Table S5-7).

3.1.2. Music perception correlates

Functional activation during music perception involved the superior and middle temporal gyri, insula, middle and inferior frontal gyri, right thalamus, hypothalamus and striatum (Fig. 1; Table S8).

3.1.3. Social cognition correlates

Functional activation during social cognition tasks involved the superior, middle and inferior temporal gyri, insula, superior, middle and inferior frontal gyri, amygdala, thalamus, striatum and anterior cingulate cortex (Fig. 1; Table S9). Repeating analyses for empathy showed activation patterns in the insulae extending to the inferior frontal gyrus, striatum, thalamus, amygdala and brainstem, the caudal temporal cortices and right postcentral gyrus (Figure S4; Table S10). Repeating analyses for Theory of Mind showed activation patterns in the bilateral temporo-parietal junction extending to the inferior frontal gyrus and precentral gyrus, precuneus, ventromedial prefrontal cortex and left striatum (Figure S4; Table S11).

3.1.4. Conjunction analyses

All conjunction analyses showed involvement of a pathway extending from the temporal cortex (BA 21, 22) through the insula (BA 13) to the inferior frontal gyri (BA 47). Additionally, basal forebrain, striatal and mesolimbic regions were activated on the right side in all conjunction analyses (Figs. 2-4; Table 4; Figure S5-6). In both FTD and

Table 1

Summary of studies included in the meta-analysis of atrophy patterns in FTD. ^aAge presented as mean \pm standard deviation or range. bvFTD = behavioral variant frontotemporal dementia; SD = semantic dementia; PNFA = progressive nonfluent aphasia; HC = healthy controls; LTLV SD = left temporal lobar variant SD; RTLW right temporal lobar variant SD; TDP = TAR DNA binding Protein 43KD, FTLD-T = frontotemporal dementia with tau inclusions; FTLD-U = frontotemporal dementia with ubiquitin inclusions; FTD-A = Alzheimer pathology with frontotemporal dementia; CDR = clinical dementia rating scale; PiD = Pick's disease; IVS 10 + 16 = mutation in exon 10 + 16; IVS 10 + 3 = mutation in exon 10 + 3; N279K = N279K mutation; S305N = S305N mutation; P301L = P301L mutation; V337M = V337M mutation. More information on the studies is presented in Table S1 of the supplementary material.

Author	Participants (%)	Age ^a	Diagnostic criteria
Agosta et al. 2009	bvFTD 31 (32%) HC 56 (58%)	58.4 \pm 10.9 66.5 \pm 9.4	Neary et al., 1998
Agosta et al. 2012	bvFTD 13 (31%) SD 7 (43%) PNFA 9 (67%) HC 25 (40%)	61.0 \pm 7.5 71.5 \pm 6.5 67.7 \pm 5.1 64.2 \pm 5.8	Neary et al., 1998
Ahmed et al. 2016	bvFTD 19 (47%) HC 25 (48%)	62 \pm 8.3 66 \pm 7.7	Rascovsky et al. 2011
Ahmed et al. 2019	bvFTD 28 (18%) HC 19 (32%)	60.9 \pm 7.0 62.9 \pm 6.9	Rascovsky et al. 2011
Ash et al. 2009	PNFA 6 SD 7 bvFTD 9 HC 31	70.7 \pm 9.3 66.8 \pm 7.3 64.8 \pm 13.2 n.a.	Neary et al., 1998
Baez et al., 2016	bvFTD 26 (46%) HC 23 (43%)	66.1 \pm 7.5 62.7 \pm 9.0	Rascovsky et al. 2011
Baez et al., 2016	bvFTD 21 (48%) HC 19 (53%)	63.8 \pm 7.3 60.4 \pm 6.8	Rascovsky et al. 2011
Baez et al. 2019	bvFTD 16 (56%) HC 22 (68%)	65.8 \pm 7.0 62.5 \pm 7.1	Rascovsky et al. 2011
Bertoux et al. 2018	bvFTD 35 HC 29	67.2 \pm 9.3 71.7 \pm 5.8	Rascovsky et al. 2011
Boccardi et al. 2005	FTD 9 (22%) HC 26 (58%)	62.0 \pm 5.0 69.0 \pm 8.0	Neary et al., 1998
Boxer et al. 2003	SD 11 HC 15	66.2 \pm 9.8 65.1 \pm 8.3	Neary et al., 1998
Brambati et al. 2009	LTLV SD 13 (69%) RTLW SD 6 (50%) HC 25 (64%)	62.0 \pm 6.3 62.5 \pm 5.8 64.8 \pm 6.9	Neary et al., 1998
Buhour et al. 2016	bvFTD 15 (67%) HC 15 (60%)	67.0 \pm 8.2 66.5 \pm 8.3	Rascovsky et al. 2011
Couto et al. 2013		69.8 \pm 7.3	Rascovsky et al. 2011; Gorno-Tempini et al., 2011

(continued on next page)

Table 1 (continued)

Author	Participants (% female)	Age ^a	Diagnostic criteria
Desgranges et al. 2007	bvFTD 12 (42%) PNFA 10 (40%) HC 18 (33%)	64.9 ± 8.6 69.8 ± 7.3	
	SD 10 HC 17	65.7 ± 8.6 65.8 ± 7.4	Neary et al., 1998
Flanagan et al. 2016	bvFTD 39 (33%) HC 61 (51%)	60.6 ± 7.5 63.6 ± 6.5	Rascovsky et al. 2011
Filippi et al. 2013	bvFTD 12 (67%) HC 30 (47%)	59.0 ± 8.0 59.0 ± 5.0	Rascovsky et al. 2011
García-Cordero et al. 2015	bvFTD 11 (55%) HC 14 (29%)	64.8 ± 5.0 57.2 ± 12.3	Rascovsky et al. 2011
Gorno-Tempini et al. 2004	PNFA 11 (73%) SD 10 (50%) HC 10 (50%)	67.9 ± 8.1 63.0 ± 5.8 69.1 ± 7.6	Neary et al., 1998
Gorno-Tempini et al. 2006	Mute PNFA 6 (83%) Nonmute PNFA 5 (100%) HC 40 (85%)	69.2 ± 8.2 62.4 ± 9.5 65.1	n.a.
Grossman et al. 2004	SD 8 PNFA 7 bvFTD 14 HC 12	65.5 ± 13.0 68.9 ± 11.4 63.1 ± 12.2 68.5 ± 9.4	Neary et al., 1998
Hardy et al. 2017	PNFA 10 (50%) SD 9 (39%) HC 19 (53%)	71.2 ± 8.9 63.8 ± 4.6 69.4 ± 4.5	Gorno-Tempini et al., 2011
Hornberger et al. 2011	bvFTD 14 (29%) HC 18 (50%)	59.3 ± 7.9 64.8 ± 5.3	Neary et al., 1998
Irish et al. 2016	SD 20 (40%) HC 35 (46%)	61.7 ± 4.8 64.4 ± 4.8	Gorno-Tempini et al., 2011
Irish et al. 2016	bvFTD 15 (40%) HC 20 (50%)	63.5 ± 7.4 67.1 ± 7.0	Rascovsky et al. 2011
Kanda et al. 2008	FTD 13 HC 20	64.9 65.2	Neary et al., 1998
Kim et al. 2007	FTLD-T 6 (17%) FTLD-U 8 (25%) HC 61 (57%)	67.7 ± 6.3 60.0 ± 10.0 68.0 ± 8.0	McKhann et al., 2001
Kipps et al. 2009	bvFTD 11 HC 12	62.1 ± 6.6 66.4 ± 4.9	Neary et al., 1998
Kumfor et al. 2018	bvFTD 19 (32%) SD 12 (50%) HC 20 (40%)	62.7 ± 8.7 64.9 ± 8.3 66.3 ± 6.1	Rascovsky et al. 2011; Gorno-Tempini et al., 2011

Table 1 (continued)

Author	Participants (% female)	Age ^a	Diagnostic criteria
Lagarde et al. 2013	bvFTD 16 (38%) HC 18 (61%)	69.3 ± 10.0 67.8 ± 5.2	Rascovsky et al. 2011
Lagarde et al. 2015	bvFTD 18 (44%) HC 18 (61%)	69.7 ± 9.7 67.8 ± 5.2	Rascovsky et al. 2011
Lee et al. 2014	bvFTD 14 (29%) HC 14 (29%)	60.8 ± 6.9 62.2 ± 4.7	Rascovsky et al. 2011
Libon et al. 2009	bvFTD 51 SD 10 PFNA 11 HC 42	61.3 ± 10.6 66.1 ± 10.8 68.7 ± 8.1 n.a.	McKhann et al., 2001; The Lund and Manchester Groups 1994
Mandelli et al. 2016	bvFTD 23 (43%) PNFA 25 (44%) HC 34 (65%)	62.9 ± 6.5 66.6 ± 7.7 62.3 ± 6.6	Rascovsky et al. 2011; Gorno-Tempini et al., 2011
Melloni et al. 2016	bvFTD 26 (46%) HC 22 (73%)	68.0 ± 11.4 68.3 ± 5.8	Rascovsky et al. 2011
Mummery et al. 2000	SD 6 (17%) HC 14 (64%)	60.5 (58–65) 62.0 (60–65)	Neary et al., 1998
Pardini et al. 2009	FTD 22 (45%) HC 14	60.3 ± 8.3 n.a.	n.a.
Pereira et al. 2009	FTD-U 9 (44%) FTD-T 6 (17%)	64.0 ± 5.7 61.8 ± 9.7	Neary et al., 1998
	PNFA 3 (0%) SD 8 (50%)	59.8 ± 7.5	
	SD-U 5 (60%)	68.3 ± 9.0	
	SD-T 3 (33%)	62.9 ± 6.4	
	FTD-A 3 (33%)	65.8 ± 6.1	
	HC 25 (44%)	58.0 ± 3.6	
Rabinovici et al. 2008	FTLD 18 (17%) HC 40 (58%)	65.3 ± 13.2 62.5 ± 8.7 63.5 ± 5.8 n.a.	McKhann et al., 2001
Rankin et al. 2011	bvFTD 5 (20%) HC 53 (55%)	63.8 ± 7.2	Cairns et al., 2007; Neary et al., 1998
Rosen et al. 2002	bvFTD 8 (25%) SD 12 (17%) HC 20 (20%)	61.8 (45–73) 67.8 (47–80) 65.4 (38–82)	Neary et al., 1998
			Neary et al., 1998

(continued on next page)

Table 1 (continued)

Author	Participants (% female)	Age ^a	Diagnostic criteria
Seeley et al. 2008	bvFTD CDR(0,5)	65.9 ±	
	15 (40%)	8.3	
	bvFTD CDR(1)	64.3 ±	
	15 (33%)	8.9	
	bvFTD CDR (2–3)	62.3 ±	
	15 (53%)	10.3	
Shen et al. 2018	HC 45 (49%)	68.3 ±	
		7.9	
	ALS-FTD 11	60.0 ±	Neary et al., 1998
	(27%)	12.7	
Whitwell et al. 2004	HC 20 (65%)	55.3 ±	
		8.4	
	FTD-T 9	52.0 ±	Neary et al., 1998
	Tau negative	8.7	
Whitwell et al. 2005	FTD 8	62.0 ±	
	HC 20	6.8	
	n.a.		
	FTD-U 9 (22%)	60.8 ±	McKhann et al., 2001
Whitwell et al. 2009	PiD 7 (43%)	2.8	
	Tau Exon 10 +	51.6 ±	
	16 5 (40%)	4.4	
	HC 20 (45%)	54.9 ±	
		2.2	
		55.7 ±	
Whitwell et al. 2011		2.8	
	IVS10 + 16 4 (25%)	56	Neary et al., 1998
	IVS10 + 3 3 (0%)	(51–62) 46 (36–49)	
	N279K 3 (100%)	49	
	S305N 2 (100%)	(43–51)	
	P301L 4 (50%)	33.5	
	V337M 3 (67%)	(34–37)	
	HC 19 (42%)	52	
		(45–65)	
		56 (49–60)	
S. M. Wilson et al. 2009	53		
		(27–65)	
	bvFTD PiD 5 (40%)	64	Neary et al., 1998
	bvFTD TDP-1 5 (80%)	68	
S. M. Wilson et al. 2010	HC 20 (50%)	63	
		(50–70)	
	SD 5 (80%)	61.4 ±	Neary et al., 1998
	HC 48 (79%)	4.8	
N. A. Wilson et al. 2020	61.5 ±		
		10.3	
	SD 25 (44%)	66.7 ±	Diagnostic guidelines
	PNFA 14 (93%)	6.0	developed by PPA researchers
	HC 10 (50%)	67.8 ±	in Buenos Aires 2006 and
		8.1	Seattle 2009
Wong et al. 2016	68.5 ±		
		5.9	
Zahn et al. 2005	bvFTD 18	n.a.	Rascovsky et al. 2011
	HC 22		
Zamboni et al. 2008	bvFTD 22 (23%)	61.0 ±	Rascovsky et al. 2011
	HC 38 (50%)	6.2	
	65.6 ±		
		5.5	
	PNFA 5 (20%)	65.0 ±	Rascovsky et al. 2011
	HC 10 (50%)	7.4	
	65.8 ±		
		7.8	
	FTD 62 (53%)	61.2 ±	Neary et al., 1998
	HC 14 (50%)	1.0	
	60.6 ±		
		1.7	

social cognition the ventromedial prefrontal cortex, the left precentral gyrus and amygdala were involved (Fig. 3; Table 4). All FTD subtypes showed overlap with both music perception and social cognition in the insulae. Whereas all FTD subtypes showed overlap with social cognition

Table 2

Summary of studies included in the meta-analysis of functional activity in music perception. ^an = number of participants. ^bAge presented as mean ± standard deviation or range. More information on the studies is presented in Table S2 of the supplementary material.

Author	Age ^b	Experimental task	Control Task
	Participants (% female) ^a		
Alluri et al. 2012	23.2 ± 3.7 n = 11 (45%)	Key clarity, increasing/decreasing pulse clarity, brightness, timbral complexity of a song 'adios nonino' by Astor Piazzolla)	
Altenmüller et al. 2014	28.7 ± 8.7 n = 18 (50%)	Participants listened to symphonic film music	Rest
Angulo-Perkins et al. 2014	28.0 ± 8.0 n = 53 (45%)	Participants listened to different musical pieces of piano, synthetic piano and violin	Non-vocal sounds and speech
Bangert et al. 2006	28.5 ± 7.3 N = 14 (57%)	Acoustic task which required passive listening to monophonic piano sequences	
Bianco et al. 2016	24.7 ± 2.9 n = 29 (59%)	Participants listened to piano pieces composed of 5 chords according to the rules of classical harmony with various melodic contours with either a congruent or incongruent ending	Rest
Bogert et al. 2016	28.2 ± 8.2 n = 56 (61%)	Participants listened to emotional musical fragments from movies	Rest
Brown and Martinez 2007	24.6 (19–46) n = 11 (55%)	1. Passive listening to piano tones 2. Melody discrimination	1. Rest 2. Button-pressing control
Chapin et al. 2010	18–29 n = 14 (64%)	Participants listened to Chopin's Etude in E major, Op.10, No. 3 was performed by a skilled pianist	Mechanical performance task
Chen et al. 2012	27.1 n = 16 (50%) (20–34)	Participants listened to melodies	
Escoffier et al. 2013	21.7 ± 1.9 n = 16 (44%)	Participants listened non-vocal musical excerpts from popular music genres	English sentences with swapped vowels
Flores-Gutiérrez et al. 2007	25.0 ± 3.1 n = 6	Participants listened to a passage of J. Prodromides, BWV 789 by J.S. Bach and G. Mahler's 5th symphony	
Groussard et al., 2010	24.6 ± 3.8 n = 20 (50%)	Participants listened to music in two parts, they then judged if these two match or are different pieces. They also judged the familiarity of musical pieces.	Sentences and familiarity of popular expressions.
Groussard et al., 2010	20–35 n = 40 (50%)	Participants judged the familiarity of musical pieces	
Habermeier et al. 2009	44.5 ± 9.9 (n = 8) n = 16 (13%) 42.9 ± 10.7 (n = 8)	Participants listened to standard and deviant melodies	

(continued on next page)

Table 2 (continued)

Author	Age ^b	Experimental task	Control Task
Participants (% female) ^a			
Janata 2009 n = 13 (85%)	20.0 (18–22)	Participants listened to music of the top 100 pop and R&B charts from when they were between 7 and 19 years of age	
Janata et al., 2002 n = 12 (58%)	28.5 (20–41)	Participants listened to different classical music parts or had to attend different elements of the music	Rest
Janata et al., 2002 n = 8 (50%)	26.0 ± 8.9	Participants listened to a melody that modulated through all 24 minor and major keys and performed tonality and timbre tasks	
Langheim et al. 2002 n = 6 (67%)	27.0 (22–32)	Participants listened to music	Rest
Levitin and Menon 2003 n = 13 (54%)	19.4–23.6	Participants listened to classical music	'Scrambled music'; quasi musical stimulus
Mitterschiffthaler et al. 2007 n = 16 (50%)	29.5 ± 5.5	Participants listened to music that was either rated as happy, sad or neutral	
Morrison et al. 2003 n = 12 (67%)	36.3	Participants (6 musicians and 6 non-musicians) listened to Western and Chinese music	Rest
Musso et al. 2015 n = 11 (55%)	24.3 (18–35)	Participants listened to chord sequences based on classical and deviant idioms	Spoken sentences which were either baseline stimuli or deviant stimuli
Park et al. 2014 n = 24 (58%)	19.0 ± 0.6 (n = 12)	Participants listened to music pieces with different emotions	Rest
Rogalsky et al. 2011 n = 20 (55%)	22.6 (18–31)	Participants listened to piano pieces	1. Rest 2. Meaningless sentences
Sachs et al. 2020 n = 40 (53%)	24.1 ± 6.2	Participants listened to sad music	
Schmitzhorst 2005 n = 15 (27%)	37.8 ± 15.2	Participants listened to famous melodies	
Toivainen et al. 2014 n = 15 (33%)	25.7 ± 5.2	Participants listened to the B-side of Abbey Road by the Beatles	
Trost et al. 2012 n = 15 (47%)	28.8 ± 9.9	Participants listened to different types of classical music from the last 4 centuries	Atonal random melodies

in striatal, basal forebrain and mesolimbic regions, only bvFTD showed overlap in these regions with music perception (Figure S7–10; Table S12–13). Subgroup analysis on empathy studies showed overlap with FTD in the insulae, the midcingulate gyrus, amygdala, striatal regions and left precentral gyrus (Figure S11–12; Table S14). Theory of Mind and FTD showed overlap in the ventromedial prefrontal cortex, temporal poles, left inferior frontal gyri connecting the left striatum (Figure S11; Table S15).

4. Discussion

In this meta-analysis we demonstrated a close neuroanatomical overlap of music perception and social cognition processing with the FTD atrophy profile.

Table 3

Summary of studies included in the meta-analysis of functional activity in social cognition. ^an = number of participants. ^bAge presented as mean ± standard deviation or range. More information on the studies is presented in Table S3 of the supplementary material. For more information on the studies see Table S3.

Author	Age ^b	Experimental task	Control task
Participants (% female) ^a			
Abraham et al. 2008 n = 17 (53%)	25.7 (22–30)	Intentional (mentalizing) relations between persons	Non-intentional relational between objects or persons
Abraham et al. 2010 n = 22 (50%)	26.1 (21–35)	Participants made judgments on intentional states based on scenarios (beliefs, desires)	Non-mental state control question
Adams et al. 2010 n = 28 (64%)	18–27	Reading the mind in the eyes task (Baron-Cohen et al. 2001)	Gender discrimination task (Baron-Cohen et al. 1999)
Aichhorn et al. 2006 n = 18 (56%)	28.5 (21–55)	Judgments about positions of objects from third person perspective	Judgments about positions of objects from first person perspective
Aichhorn et al. 2009 n = 21 (38%)	24 (21–41)	False belief stories	False photograph stories
Akitsuki and Decety 2009 n = 26 (54%)	24.4 ± 5.8	Participants watched painful stimuli	Non-painful stimuli
Assaf et al. 2009 n = 19 (47%)	32.3 ± 10.4	Participants play a game where mentalizing is needed	A choice in the game where no mentalizing is needed
Azevedo et al. 2013 n = 27 (59%)	23.9	Participants watched painful stimuli	Non-painful stimuli
Azevedo et al. 2014 n = 12 (100%)	22.2 ± 2.6	Participants watched painful stimuli	Non-painful stimuli
Bahnemann et al. 2009 n = 25 (12%)	26 ± 4.4	Judgment of mental states from a movie (Dziobek et al. 2006)	Participants make appearance judgments in the movie (Baron-Cohen et al. 1999)
Benuzzi et al. 2008 n = 15 (100%)	23.5 (19–31)	Participants watched painful stimuli	Non-painful stimuli
Benuzzi et al. 2018 n = 27 (100%)	21.3 ± 1.6	Participants watched painful facial expressions	Neutral facial expressions
Berlingeri et al. 2016 n = 25 (52%)	25.3 ± 4.8	Participants judged painful experience of a character	Non-painful stimuli
Boccadoro et al. 2019 n = 68 (75%)	31.1 ± 10.5	Participants watched false belief videos to activate spontaneous ToM	True belief videos
Bodden et al. 2013 n = 30 (50%)	25.3 ± 2.5	Judgment about the affective and cognitive aspects of a scenario	Participants answered questions about the physical state of a scenario
Bos et al. 2015 n = 24 (0%)	23.1 (19–27)	Participants watched painful stimuli	Non-painful stimuli
Botvinick et al. 2005 n = 12 (100%)	20–30	Participants watched painful facial expressions	Neutral facial expressions
Brüne et al. 2008 n = 13 (70%)	26.5 ± 5.3	Participants judged intentions and expectations of the protagonist in a cartoon	Participants properties of objects in the cartoon
Bruneau et al. 2012 n = 14 (57%)	23.5 ± 4.1	Participants read stories about painful physical and emotional scenarios	Non painful physical and emotional scenarios

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Brunneau et al. 2015 n = 18 (78%)	22.2 ± 3.6	Participants actively emphasized in emotional painful scenarios	Participants stayed objective
Brunnlieb et al. 2013 n = 18 (0%)	19–37	Participants watched pictures of an emotionally charged situation (Krämer et al. 2010)	Pictures of neutral situations
Budell et al. 2010 n = 18 (50%)	18–25	Participants judged the amount of pain in painful facial expressions	Neutral facial expressions
Canessa et al. 2012 n = 27 (52%)	26.3 ± 4.2	Participants viewed images displaying affective social interactions	Images displaying natural or urban landscapes
Cassidy et al. 2020 n = 75 (63%)	21.6 ± 2.8 71.7 ± 6.1 (n = 35)	False belief stories (Saxe and Kanwisher 2003)	False photographs
Castelli et al. 2010 n = 24 (75%)	25.2 ± 3.5 (n = 12) 65.2 ± 5.7 (n = 12)	Reading the mind in the eyes task (Baron-Cohen et al. 2001)	Gender discrimination task (Baron-Cohen et al. 1999)
Chaminade et al. 2012 n = 19 (0%)	21.5 ± 4.9	A competitive game was played against a fellow human	A computer playing randomly
Cheng et al. 2007 n = 14 (50%)	n.a.	Participants watched painful stimuli	Non-painful stimuli
Cheng et al. 2010 n = 36 (50%)	23 ± 3	Participants watched painful stimuli	Non-painful stimuli
Cheon et al. 2013 n = 27 (44%)	23.1 ± 4.4 (n = 13) 25.1 ± 4.8 (n = 14)	Participants watched pictures of emotionally painful scenarios	Non-painful scenarios
Cheung et al. 2012 n = 23	23.5 ± 1.1	Cartoon false belief stories	Non-mentalistic pictures and verbal stories (fillers)
Chiao et al. 2009 n = 14 (100%)	22.9 ± 3.7	Participants watched pictures of emotionally painful scenarios	Non-painful scenarios
Christov-Moore and Iacoboni 2019 n = 70 (51%)	18–35	Participants watched painful stimuli	Non-painful stimuli
Contreras et al. 2013 n = 14 (43%)	19.9 (18–23)	False belief stories (Saxe and Kanwisher 2003)	False photographs
Corradi-Dell'Acqua et al. 2011 n = 28 (100%)	19–31	Participants watched painful noxious stimuli	Non-painful/noxious stimuli
Corradi-Dell'Acqua et al. 2014 n = 46 (100%)	18–31	Judgment about a protagonists beliefs, emotions and pain (Saxe and Kanwisher 2003)	Stories without a protagonist, but with physical representation on a map or photograph
Danziger et al. 2009 n = 13 (54%)	33 ± 9.0	Participants watched painful stimuli and facial expressions	Non-painful stimuli and neutral expressions
de Achával et al. 2012 n = 14 (43%)	28.4 ± 8.3	Reading the mind in the eyes task (Saxe and Kanwisher 2003)	Gender discrimination task (Baron-Cohen et al. 1999)
de Grecq et al., 2012 n = 20 (60%)	37.0 ± 10.6	Participants actively emphasized with characters	Smoothed pictures
Grecq et al., 2012 n = 20 (55%)	23 (21–26)	Participants actively emphasized with characters	Skin color evaluation
	27.0 ± 5.0		Neutral expressions

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Deeley et al. 2006 n = 9 (0%)	28.3 ± 6.6 (n = 12) 26.3 ± 4.1 (n = 12)	Participants watched fearful facial expressions	1. Participants judged emotions from faces 2. Emotional perspective taking task (ToM) 3. Affective responsiveness task (empathy)
Derntl et al. 2010 n = 24 (50%)	22.0 ± 3.2	1. Participants judged emotions from faces 2. Emotional perspective taking task (ToM) 3. Affective responsiveness task (empathy)	1. Gender discrimination task 2. Characteristics task 3. Sentence forming task
Deuse et al. 2016 n = 38 (53%)	24.1 (18–30)	Mental state judgments were made about daily life situations as pleasant or unpleasant on a 9-point scale	Comparable scenes to the experimental task but without people
Dodell-Feder et al. 2011 n = 62 (65%)	24.7 ± 2.2	False belief story "Sally Anne paradigm" (Baron-Cohen et al. 1985)	Non-mental control questions
Döhnel et al. 2012 n = 18 (44%)	25.3 ± 4.6	Emotional and behavioral "Sally Anne paradigm" (Baron-Cohen et al. 1985)	True belief stories
Dufour et al. 2013 n = 462 (52%)	24.9 (18–69)	False belief stories	False photographs
Dungan and Young 2019 n = 26 (52%)	23.7 ± 4.4	Participants judged mental states (why)	Participants judged behavior (how)
Enzi et al. 2016 n = 20 (0%)	27.0 ± 5.1	Participants watched painful stimuli	Non-painful stimuli
Ernst et al. 2013 n = 18 (67%)	27.0 ± 7.6	Participants judged the empathic ability to faces	Smoothed photographs
Fan et al. 2014 n = 21 (0%)	19.3 ± 3.4	Participants watched painful stimuli	Non-painful stimuli
Frank et al. 2015 n = 34 (50%)	28 ± 0.42 (n = 17) 29 ± 0.5 (n = 17)	False belief stories	Unlinked sentences
Feng et al. 2016 n = 22 (50%)	22.2 ± 1.9	Participants watched painful stimuli	Non-painful stimuli
Fourie et al. 2017 n = 38 (55%)	40.1 ± 4.1 (n = 19) 41.5 ± 5.8 (n = 19)	Participants watched facial expressions of people in physical pain and scenarios of social pain	Neutral expressions and scenarios
Fujino et al. 2014 n = 11 (82%)	32.3 ± 10.5	Participants watched painful stimuli	Non-painful stimuli
Gallagher et al. 2000 n = 6 (17%)	30.0 (23–36)	False belief stories (Fletcher et al. 1995)	Participants judged why something physical happened in a non-ToM story
Gallagher and Frith 2004 n = 12 (42%)	36.0 (28.4–59.5)	Participants judged expression of inner states by gestures	Neutral gestures
Geiger et al. 2019 n = 32 (47%)	25.8 ± 4.9	Participants judged emotional states from animations	Judgment of activities
Gilbert et al. 2007 n = 16 (56%)	21 (18–27)	Participants judged if the experimenter was trying to be helpful in a task of time	Participants are told a computer manages time randomly
Gobbini et al. 2007 n = 12 (58%)	22.0 ± 2.0	False belief stories (Fletcher et al. 1995)	Questions about stories describing human activity

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
		Participants (% female) ^a	
Göttlich et al. 2017	27.8 ± 4.8 (n = 17)	Participants viewed socio-affective scenarios to elicit empathetic responses	without mental state attribution Neutral scenarios
n = 44 (75%)	23.0 ± 3.3 (n = 27)		
Grèzes et al. 2007	25	Participants watched fearful body expressions	Neutral body expressions
n = 16 (63%)			
Grice-Jackson et al. 2017	24.0 ± 1.4	Participants judged whether they experienced pain while watching painful stimuli	Non-painful stimuli
n = 44 (59%)			
Gu and Han 2007	21.0 ± 1.2	Participants watched painful stimuli	Non-painful stimuli
n = 12 (42%)			
Gu et al. 2010	24.8	Participants made pain judgments from painful stimuli	Non-painful stimuli
n = 18 (50%)	(22–28)		
Gu et al. 2013	25.2	Participants made pain judgments from painful stimuli	Non-painful stimuli
n = 18 (50%)	(22–34)		
Guo et al. 2013	22.2 ± 2.7	Participants watched painful stimuli	Non-painful stimuli
n = 40 (75%)			
Gweon et al. 2012	21.5	Stories about a protagonists mental state	Stories about physical events
n = 8 (75%)	(18–25)		
Haas et al. 2015	21.8 ± 2.1	An emotion attribution task judging the cause of an emotional response	Gender match task
n = 16 (56%)			
Hadjikhani et al. 2014	22.5 ± 7.5	Participants watched painful facial expressions	Non-painful stimuli
n = 31 (10%)			
Han et al. 2009	21.0 ± 1.6	Participants watched painful stimuli with or without painful expressions	Non-painful stimuli and neutral expressions
n = 46 (50%)	(n = 24)		
	22.6 ± 2.3 (n = 22)		
Han et al. 2017)	22.9	Participants watched painful stimuli	Non-painful stimuli
n = 33 (48%)			
Harris et al. 2005	20	Participants judged behavior to the characteristics of a person	Baseline condition
n = 12 (92%)			
Hervé et al. 2013	30.9 ± 8.6	Participants judged the mental state of a character	Participants judged if a sentence was true or not
n = 42 (38%)			
Hooker et al. 2008	21.0	False belief stories (Fletcher et al. 1995)	True belief stories
n = 20 (55%)	(19–26)		
Hooker et al. 2010	21 (18–25)	Participants viewed a scenario with change in beliefs and mental states	No change in beliefs and mental states
n = 15 (47%)			
Jackson et al. 2005	22 ± 2.6	Participants watched painful stimuli	Non-painful stimuli
n = 15 (47%)			
Jackson et al. 2006	29 ± 6.5	Participants watched painful stimuli	Non-painful stimuli
n = 34 (59%)			
Jacoby et al. 2016	25.3	1. False belief task (ToM; Dodell-Feder et al. 2011) 2. Stories about painful physical and emotional scenarios (empathy; Bruneau et al. 2012)	1. False photographs or maps 2. Non-painful physical and emotional scenarios
n = 20 (60%)	(18–39)		
Jankowiak-Siuda et al. 2015	25–35	Participants watched painful stimuli	Non-painful stimuli
n = 27 (52%)			

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
		Participants (% female) ^a	
Jenkins and Mitchell 2010	19.8 (18–22)	Participants judged mental states in an (un) ambiguous scenario	Nonsocial physical representations (e.g. photographs)
n = 15 (60%)			
Jimura et al. 2010	20–28	False belief stories (Saxe and Kanwisher 2003)	Participants judged physical aspects of the story
n = 34 (47%)			
Kaiser et al. 2008	28.7 ± 6.6 (n = 12)	Participants judged a characters visual perspective	Own perspective
n = 24 (50%)	27.6 ± 4.9 (n = 12)		
Kana et al. 2009	24.4 ± 3.7 (n = 12 (0%))	Mental states attribution to movement of geometrical figures (Castelli et al. 2002)	Random movements
Kana et al. 2016	24.0 ± 13.5 (n = 15)	Participants judged emotional facial/ body expressions	Neutral expressions
Kana and Travers 2012	21.0 (18.5–35.8)	Participants judged emotional body expressions	Participants judged actions
n = 26 (54%)			
Kandylaki et al. 2015	24.3 ± 2.1 (n = 20)	Participants listened to stories with false belief passages and judged beliefs	Judgments about physical events in the story.
Kanske et al. 2015	40.9 ± 9.5 (n = 178 (60%))	1. False belief task (ToM; Dodell-Feder et al. 2011) 2. Socio-affective video task (empathy; Klimecki et al. 2013)	1. nonToM story contents 2. Neutral videos
Kim et al. 2005	23.3 ± 2.0 (n = 14 (50%))	Participant judged if the facial affect was appropriate for a situation	Participant were asked whether the gender of two subjects was the same
Kim et al. 2007	25.4 ± 2.6 (n = 21 (48%))	Participants imagined facial affective expression of emotional faces	Imagery of neutral faces
Kim et al. 2009	27.5 ± 3.3 (n = 14 (43%))	Participants made judgments on causality of facial expressions	Clear information on causality
Kircher et al. 2009	27.4 (n = 14 (0%))	Participants played a mentalizing game	Non-mentalizing game
Kobayashi et al. 2008	29.7 ± 4.6 (n = 16 (50%))	False belief stories	Baseline condition
Kockler et al. 2010	24.4 ± 2.1 (n = 18 (0%))	Participants judged a characters visual perspective	Own perspective
Koelkebeck et al. 2011	30.9 ± 8.1 (n = 15 (53%))	Judgment of 'moving shapes' paradigm with interacting triangles (Abell et al. 2000)	Random movements
Krach et al. 2008	24.5 ± 3.0 (n = 20 (0%))	Participants played a mentalizing game ('Prisoners dilemma game')	Control condition
Krach et al. 2009	27.4 (n = 24 (50%))	Participants played a mentalizing game ('Prisoners dilemma game')	Baseline game
Krach et al. 2011	22.8 ± 2.2 (n = 32 (53%))	Participants watched vicarious embarrassing social situations	Neutral situations
Krach et al. 2015	24.3 ± 2.9 (n = 16 (0%))	Participants watched vicarious embarrassing social	Neutral situations and non-painful stimuli

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
		Participants (%) female) ^a	
Krämer et al. 2010 n = 16 (63%)	27.8 ± 4.8	situations and painful stimuli Participants were showed pictures of an emotionally charged situation	Pictures of neutral situations
Lamm and Decety 2008 n = 18 (50%)	23.7 ± 4.0	Participants watched painful stimuli	Non-painful stimuli
Lancaster et al. 2015 n = 37 (0%)	23.7 (18–29)	Social attribution task (Schultz et al. 2003)	Random movements
Lavoie et al. 2016 n = 19 (16%)	28.8 ± 7.9	Participants judged a characters feelings or thoughts	Participants answered questions about physical events in a scenario Physical properties
Lawrence et al. 2006 n = 12 (50%)	32.2 ± 9.9	Participants performed emotion perception tasks	
Lee et al. 2010 n = 18 (50%)	25.8 ± 2.2	Participants judged comics on various types of empathic causality	Physical causality
Lee et al. 2013 n = 12 (100%)	72.3 ± 6.2	Participants watched emotional facial expressions	Neutral expressions
Leiberg et al. 2012 n = 24 (100%)	24.1 (18–33)	Participants viewed emotional scenarios	Neutral scenarios
Leitman et al. 2010 n = 20 (0%)	28 ± 5	Participants judged emotional prosody from voices	Neutral prosody judgment
Lewis et al. 2017 n = 17 (53%)	22 ± 2.9	Participants answered true/false mentalizing questions	Factual memory processing
Liew et al. 2011 n = 18 (44%)	23 ± 2.3	Participants judged intentions of actors using gestures	Still photo of the actor
Lin et al. 2018 n = 39 (46%)	22.2 ± 2.7	False belief task (Dodell-Feder et al. 2011)	False photo task
Lombardo et al. 2010 n = 33 (0%)	28.0 ± 6.1	Participants answered questions about others mental states	Participants answered questions about physical characteristics
Lotze et al. 2006 n = 20 (50%)	52.3 ± 12.4	Participants viewed emotional/expressive gestures	Isolated hand movements
Luo et al. 2014 n = 36 (50%)	21.3	Participants watched painful stimuli	Non-painful stimuli
Malhi et al. 2008 n = 20 (45%)	35.8 ± 10.4	Observing geometric shapes interacting (Castelli et al. 2000)	Random movements
Mano et al. 2009 n = 18 (56%)	25.2 ± 2.5	Participants judged emotions and beliefs in a scenario	Unrelated stories
Marjoram et al. 2006 n = 13 (38%)	29.6 ± 1.6	Participants viewed jokes needing ToM for understanding (Gallagher et al. 2000)	Physical jokes
Martin and Weisberg 2003 n = 12 (50%)	27.5 (23–34)	Geometric shapes conveying social interactions	Mechanical actions
Mathur et al. 2010 n = 28 (46%)	23.8 ± 0.8	Participants watched people in emotionally painful situations	Neutral situations
Mathur et al. 2016 n = 15 (53%)	25.3 ± 4.7	Participants watched people, animals and nature in emotionally painful situations	Neutral situations
Mazza et al. 2013 n = 10 (100%)	24.4 ± 4.4	Participants judged pictures with negative emotional valence	Neutral pictures

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
		Participants (%) female) ^a	
Mazzola et al. 2010 n = 30 (37%)	36.8	Participants watched painful facial expressions	Neutral facial expressions
McAdams and Krawczyk 2011 n = 17 (100%)	24.7 ± 6.6	Participants judged how shapes interacted	Visuospatial task
Meaux and Vuilleumier 2016 n = 26 (50%)	25.9 ± 5.5	Participants judged facial emotions (happy/angry)	Neutral emotions
Melchers et al. 2015 n = 60 (65%)	29.5 ± 5.4	Participants watched vicarious embarrassing social situations	Neutral situations
Mercadillo et al. 2011 n = 24 (50%)	27.0 ± 2.5	Participants watched social scenarios of people suffering	Neutral social scenarios
Meyer et al. 2015 n = 25 (60%)	21.6 ± 2.5	Participants made personality trait judgments about friends	Participants alphabetized the names
Mier et al., 2010 n = 40 (50%)	25.3 ± 3.5	Participants made emotion recognition and mentalizing judgments from faces	Judgment of physical features
Mier et al., 2010 n = 16 (31%)	37.0 ± 8.2	Participants made emotion recognition and mentalizing judgments from faces	Judgment of physical features
Mitchell 2008 n = 20 (55%)	23 (19–29)	False belief stories (Saxe and Kanwisher 2003)	False photograph stories
Moessnang et al. 2017 n = 28 (50%)	22.9 ± 2.8	Judgment of 'moving shapes' paradigm with interacting triangles (Abell et al. 2000)	Goal directed movements
Moll et al. 2007 n = 12 (50%)	28.5 ± 9.6	Participants read scenarios that evoked empathetic/compassionate feelings	Neutral scenarios
Moran et al. 2012 n = 31	23.0 ± 0.9	Animate movement task, moral judgment task (Young et al. 2007) and False belief stories (Saxe and Kanwisher 2003)	Mechanical movies, neutral outcome and false photo stories
Morelli et al. 2014 n = 32 (50%)	19.9 ± 1.4	Participants empathized with people in photo's of emotional scenarios	Neutral images
Moriguchi et al. 2007 n = 14 (86%)	20.8 ± 0.9	Participants watched painful stimuli (Jackson et al. 2005)	Non-painful stimuli
Morrison et al. 2004 n = 14 (64%)	23.0	Participants watched painful stimuli	Non-painful stimuli
Morrison and Downing 2007 n = 12 (42%)	31.0	Participants watched painful stimuli	Non-painful stimuli
Morrison et al. 2007 n = 16 (50%)	27.0	Participants watched painful stimuli	Non-painful stimuli
Morrison et al. 2013 n = 14 (50%)	23–35	Participants watched painful stimuli	Non-painful stimuli
Murphy et al. 2010 n = 10 (60%)	29.6 ± 8.4	Participants judged personality attributes	Semantic positivity evaluation of adjectives
Narumoto et al. 2000 n = 8 (38%)	23–29	Facial emotion recognition task	Gender matching task

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Nomi et al. 2008 n = 14 (50%)	28.6 ± 5.5	Participants recognized and shared emotions of presented facial expressions	Count earrings
Nummenmaa et al. 2008 n = 10 (100%)	26 ± 5.6	Participants empathized with emotional scenes	Neutral scenes
Ochsner et al. 2004 n = 13 (54%) ⁷	29.5	Participants judged whether a character felt pleasant, unpleasant or neutral	Participants judged whether a photo was taken outside or inside
Otti et al. 2015 n = 20 (65%)	45.6 ± 14.0	Participants judged a scenario where triangles were coaxing, mocking, seducing and surprising each other	Participants judged a scenario where triangles were moving or rotating without any interaction
Özdem et al. 2017 n = 20	18–26	False belief stories (Saxe and Kanwisher 2003)	False photograph stories
Paulus et al. 2018 n = 17 (0%)	24.1 ± 3.0	Participants emphasized with characters in embarrassing scenarios	Neutral scenarios
Pichon et al. 2008 n = 16 (44%)	18–26	Participants judged emotions from angry whole body expression	Neutral body expressions
Platek et al. 2004 n = 5	n.a.	Reading the mind in the eyes task (Baron-Cohen et al. 2001)	Participants viewed a crosshair
Powell et al. 2017 n = 12 (0%)	36.4 ± 13.9	Intentional judgments and mental state attributions (ToM & empathy; Brunet et al. 2000)	Physical causality
Preis et al. 2013 n = 64 (50%)	22.9	Participants watched painful stimuli	Non-painful stimuli
Prochnow et al. 2013 n = 15 (47%)	22.3 ± 1.4	Participants watched and judged emotional faces and gestures	Scrambled stimuli
Prochnow et al. 2014 n = 26 (47%)	38.3 ± 12.0	Participants matched facial expressions with situations	Scrambled images with unrelated sentences
Qiao-Tassert et al. 2018 n = 24 (54%)	27.6 ± 6.0	Participants watched painful stimuli	Non-painful stimuli
Rabin et al. 2010 n = 18 (50%)	57.2 ± 8.0	Participants viewed photo's and judged mental state attributions vividly	Scenarios lacking details
Rapp et al. 2010 n = 15 (100%)	28.1 ± 8.0	Participants judged irony of sentences	Literal target sentences
Regenbogen et al. 2012 n = 27 (48%)	34.1 ± 9.8	Participants judged valence from emotional prosody, facial expressions and speech content	Neutral content
Reniers et al. 2014 n = 15 (0%)	18–40	Imagining what someone feels (empathy) and would make them feel better (ToM)	Neutral contexts
Rilling et al. 2008 n = 20 (100%)	20.8 ± 1.6	Participants played a mentalizing game ('Prisoners dilemma game')	Gamble game
Rosenblau et al. 2016 n = 22 (27%)	31.3 ± 8.5	Participants judged changes in affective	Physical inference tasks

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Roser et al. 2012 n = 14 (0%)	27.3 ± 3.5	states of a protagonist in a video	Participants judged intentions and beliefs in cartoons (Brüne 2005)
Rothmayr et al. 2011 n = 12 (58%)	23.7 (23–24)	False belief story ''Sally Anne paradigm'' (Baron-Cohen et al. 1985).	True belief story
Ruckmann et al. 2015 n = 30 (50%)	24.5 ± 3.7	Participants watched painful stimuli	Non-painful stimuli
Saft et al. 2013 n = 26 (27%)	28.8 ± 4.1	Participants judged intentions and beliefs in cartoons (Martin Brüne 2005)	Participants judged properties of objects displayed
Samson et al. 2008 n = 17 (53%)	26.1 ± 3.3	Participants viewed comic cartoons that required ToM for understanding	Non-ToM cartoons
Saxe and Powell 2006 n = 12 (75%)	19–26	1. False belief stories 2. 'Thought stories' where participants judged protagonist's beliefs and thoughts	1. False-photograph stories 2. A story of the protagonist's physical, social characteristics and physical feelings Non-mentalizing questions about the stories
Schlaffke et al. 2015 n = 39 (0%)	25.9 ± 5.8	Participants judged a characters feelings (affective ToM) and beliefs (cognitive ToM)	Non-mentalizing questions about the stories
Schmitgen et al. 2016 n = 21 (9%)	23.7 ± 3.1	Participants judged a protagonist's mental state as worse, equal or better after an event	Participants judged the number of living beings in a scene
Schnell et al. 2011 n = 21 (43%)	25.5 ± 5.0	Participants viewed cartoons, which they judged affective and visuospatial states	Participants answered questions about their own affective and visuospatial states
Schulte-Rüther et al. 2008 n = 26 (54%)	24.8 ± 3.7	Emotion recognition and affect task from facial expressions	Age/gender task
Schuwerk et al. 2017 n = 24 (50%)	22.8 ± 3.0	Participants belief was they received cues from a person	Participants belief was they received cues from the computer
Seara-Cardoso et al. 2015 n = 46 (0%)	27.9 (19–40)	Participants watched painful stimuli	Non-painful stimuli
Seara-Cardoso et al. 2016 n = 30 (0%)	26.9 (20–40)	Participants judged their own affective state watching facial expressions	Fixation cross
Seger et al. 2004 n = 12 (67%)	23.6 (20–32)	Participants made judgments if someone would like particular food	Judgments about the number of vowels
Sheng et al. 2014 n = 21 (52%)	22.0 ± 1.8	Participants judged painful facial expressions	Neutral facial expressions
Shibata et al. 2010 n = 13 (23%)	23.8 (20–29)	Participants judged irony of sentences	Literal sentences
Shibata et al. 2011 n = 15 (27%)	25.2	Participants judged the meaning of indirect sentences	Literal sentences
Simon et al. 2006 n = 17 (53%)	23.1 ± 4.1	Participants judged gender on painful facial expressions	Neutral facial expressions
	25.8 ± 6.9		Non-painful stimuli

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Singer et al. 2004 n = 16 (100%)		Participants watched painful stimuli	
Sommer et al. 2007 n = 16 (50%)	26.0 (23–37)	False belief story "Sally Anne paradigm" (Simon Baron-Cohen et al. 1985)	True belief story
Specht and Wigglesworth 2018 n = 18 (0%)	25.7 ± 2.5	Participants judged intentions of a character (Brunet et al. 2000)	Participants judged which cartoon was displayed twice
Spiers and Maguire 2006 n = 20 (0%)	49.8 ± 8.5	Mental state attributions during a game	Coasting condition
Spotorno et al. 2012 n = 20 (60%)	22	Participants judged irony of sentences	Literal sentences
Sprengelmeyer et al. 1998 n = 6 (67%)	23.5 ± 1.3	Participants judged gender on emotional facial expressions	Neutral facial expressions
Spunt et al. 2011 n = 15 (60%)	19.5 ± 1.9	Participants judged the motive of a character ('why')	Participants judged what a character was doing
Spunt and Ralph Adolphs, 2014 n = 29 (34%)	27.1 (19–38)	1. False belief stories (Dodell-Feder et al. 2011; Saxe and Kanwisher 2003) 2. Participants judged motive	1. False photograph stories 2. Participants judged physical events
Spunt and Lieberman 2012 n = 22 (55%)	21.6 (19–32)	Participants judged the motive of a character ('why')	Participants judged what a character was doing
Spunt et al. 2017 n = 16 (50%)	29.0 (21–46)	Participants judged an emotion from a human, dog or monkey	Participants judged characteristics of facial expressions
Sripada et al. 2009 n = 26 (62%)	28.0 ± 8.2	Participants played the 'trust game' against a human	Participants played against a computer
Suzuki et al. 2012 n = 26	20–35	Participants predicted the choices of another person	Random rewards
Takahashi et al. 2015 n = 38 (100%)	22.1 ± 4.7	Participants judged the intensity of sadness of faces with tears	Participants judged the intensity of sadness of faces without tears
Tamm et al. 2017 n = 86 (51%)	23.0 (n = 47) 68.0 (n = 39)	Participants watched painful stimuli and rated their own unpleasant affect	Non-painful stimuli
Tholen et al. 2020 n = 130 (55%)	40.4 ± 9.0	1. False belief task (ToM; Dodell-Feder et al. 2011) 2. Socio-affective video task (empathy; Klimecki et al. 2013)	1. nonToM story contents 2. Neutral videos
Thye et al. 2018 18 = (72%)	20.2 ± 1.4	1. Reading the mind in the eyes (S Baron-Cohen et al. 2001) 2. Reading the mind in the voice (Rutherford, 2002) 3. Intentional judgments (Brunet et al. 2000)	1. Gender discrimination task 2. Gender discrimination task 3. Physical events
Todorov et al. 2007 n = 11 (27%)	21 (18–32)	Participants judged faces based on their personality traits	Novel faces
Uchiyama et al. 2006 n = 20 (50%)	21.9 ± 2.7	Participants judged if a sentence expressed sarcasm	Literal sentences
	27.3		Static right hand

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (%) female) ^a			
Ushida et al. 2008 n = 15 (53%)		Participants watched painful stimuli	
Vachon-Presseau et al. 2012 n = 20 (50%)	36 ± 10	Participants watched painful stimuli and painful facial expressions	Neutral stimuli
van Ackeren et al. 2016 n = 25 (100%)	18–35	Participants judged if a sentence expressed indirect replies and requests	Direct replies
van der Meer et al. 2011 n = 19 (47%)	21.6 ± 2.6	False belief stories	Physical events in a scenario
Van Hoeck et al. 2014 n = 19	19–29	False belief stories	True belief
Vanderwal et al. 2008 n = 17 (41%)	21.5 ± 1.8	Social attribution task (Schultz et al. 2003)	Bumper cars condition
Veroude et al. 2012 n = 25 (100%)	19.1 (18.1–19.9)	Participants made mental state judgments	Baseline condition
Vistoli et al. 2016 n = 21 (14%)	29.2 ± 7.9	Participants watched painful stimuli	Non-painful stimuli
Vogeley et al. 2001 n = 8 (0%)	25–36	Participants judged the mental state of a character (Fletcher et al. 1995)	Sentences with no semantic consistency or coherence
Völlm et al. 2006 n = 13 (0%)	24.9 (19–36)	Intentional judgments (Brunet et al. 2000) and empathic judgment	Physical causality
Walter et al. 2004 n = 13 (54%)	25.2 ± 2.0	Participants judged communicative intentions in stories	Physical causality
Walter et al. 2009 n = 12 (50%)	24.8 ± 2.6	Participants judged private and social intentions in stories	Physical causality
Wang et al. 2006 n = 12 (50%)	26.9 ± 3.5	Participants judged irony of sentences	Literal sentences
Wang et al. 2015 n = 56 (45%)	19.3 ± 0.9	Participants performed ToM and empathy tasks (Völlm et al. 2006)	Physical causality
Whitehead and Armony 2019 n = 30 (50%)	24.0 ± 2.6	Participants judged expressions of fear from faces, bodies, prosody and vocalization	Neutral expressions
Wicker et al. 2003 n = 14 (0%)	20–27	Participants watched facial expressions of disgust and pleasure	Neutral expressions
Wolf et al. 2010 n = 18	20–45	Judgment of mental states from a movie (Dziobek et al. 2006)	Physical details
Young et al. 2007 n = 27 (44%)	18–22	Participants judged belief states of a protagonist	False photograph stories
Young et al. 2010 n = 17 (41%)	18–31	False belief stories	False photograph stories
Young et al. 2011 n = 17 (59%)	18–22	False belief stories (Saxe and Kanwisher 2003)	False photographs or maps
Zaitchik et al. 2010 n = 15 (47%)	22.4 (20–24)	Participants judged beliefs and emotional states	Syntax control condition
Zhang et al. 2018 n = 25 (52%)	20.9 (19–24)	Participants judged emotions from facial expressions and vocal prosody	Neutral faces and prosody

(continued on next page)

Table 3 (continued)

Author	Age ^b	Experimental task	Control task
Participants (% female) ^a			
Zheng et al., 2016 n = 20 (50%)	25.0 ± 1.6	Participants watched painful stimuli	Non-painful stimuli
Zheng et al., 2016 n = 20 (60%)	21.7 ± 1.9	Participants watched painful stimuli	Non-painful stimuli
Ziae et al. 2016 n = 40 69.8 ± 3.0 (n = 20)	20.7 ± 2.7 (n = 20)	Reading the mind in the eyes task (Baron-Cohen et al. 2001)	Gender discrimination task




Fig. 1. Meta-analytic results of brain regions involved in atrophy of FTD and functional activity during music perception and social cognition tasks. z = axial location of x,y,z coordinates. x = sagittal location of x,y,z coordinates. The ALE-scores are demonstrated. For the coordinates and brain regions see Table S4, S8-9. All results were thresholded at cluster-wise threshold p < 0.05 (FWE-corrected).

Our results were in line with previous research for the observed atrophy patterns in FTD (C. Luo et al. 2020; Pan et al. 2012; Schroeter et al. 2007, 2008; Yang et al. 2012) and functional activity in music perception (Janata 2015; Limb 2006). The observed functional activity pattern in social cognition was similar to previous meta-analyses on Theory of Mind (Molenberghs et al. 2016; Schurz et al. 2014). Unlike these studies, we only observed activation in the precuneus in the Theory of Mind subgroup analysis, but not general social cognition. This suggests that activation in the precuneus might be specific for Theory of Mind tasks.

Our study extends the literature by demonstrating evidence for a potential relationship between the three modalities. We found that neuronal circuits showing atrophy in FTD patients are also functionally involved in music perception and social cognition. This provides a putative biological substrate for alterations in music perception and social cognition observed clinically. The shared anatomical areas in music perception and social cognition extended from the caudal part of the superior temporal gyrus (BA 21) rostrally to the inferior frontal gyrus (BA 47). This neuronal circuit was also observed in the conjunction analysis of FTD and social cognition and thus appears to play a central role in all three modalities. In the left hemisphere these brain areas are part of the ‘ventral language pathway’ that extends from the auditory cortex through the insula to the inferior frontal gyrus (Saur et al. 2008), and is known to be involved in higher-level language processing having mainly a sound-to-meaning function (Aryani et al. 2019; Berwick et al. 2013; Fröhholz et al. 2016; Saur et al. 2008). Atrophy in most of these




Fig. 2. Brain regions involved in the conjunction analysis of FTD and music perception. z = axial location of x,y,z coordinates. y = coronal location of x,y,z coordinates. The ALE-scores are demonstrated. For the coordinates and brain regions see Table 4. All results were thresholded at a cluster-wise threshold of p < 0.05 (FWE-corrected).




Fig. 3. Brain regions involved in the conjunction analysis of FTD and social cognition. y = coronal location of x,y,z coordinates. The ALE-scores are demonstrated. For the coordinates and brain regions see Table 4. All results were thresholded at a cluster-wise threshold of p < 0.05 (FWE-corrected).

brain areas were previously reported to be related to a loss of sarcasm recognition as part of social cognition tasks in FTD (Downey et al. 2015). Oechslin et al. reported that the ventral pathway in the right hemisphere




Fig. 4. Brain regions involved in the conjunction analysis of music perception and social cognition. z = axial location of x,y,z coordinates. y = coronal location of x,y,z coordinates. The ALE-scores are demonstrated. For the coordinates and brain regions see Table 4. All results were thresholded at a cluster-wise threshold of $p < 0.05$ (FWE-corrected).

was functionally involved in music syntax processing in musicians (Oechslin et al. 2017). Other studies found similar brain regions involved in affective voice processing (Leitman et al. 2010; Regenbogen et al. 2012; Zhang et al. 2018). Together with our results, this suggests that brain regions involved in interpreting sounds of social cognition and music may share a neurobiological basis bilaterally, and could explain loss of these functions in FTD. Of note is that the studies we included in this meta-analysis for social cognition used different test paradigms, which did not necessarily aim at the interpretation of voices. The observation that this pathway was robustly implied with social cognition suggests that it might be a key circuit for social cognition. All three modalities further demonstrated an anatomical overlap in mesolimbic dopaminergic circuits which are also known to be involved in reward and behavior (Omar et al. 2011; Salimpoor et al. 2011; Zatorre and Salimpoor 2013) and basal forebrain and striatal regions that are part of the semantic appraisal network which is involved in weighing hedonic value of (social and asocial) stimuli (Ranasinghe et al. 2016; Seeley et al. 2012; Zhou and Seeley 2014). These findings possibly explain the altered rewarding sensation that patients with FTD can experience when listening to music (i.e. musicophilia or sound-aversion; Fletcher et al. 2013; Omar et al. 2011). Of note, in the subgroup analyses we only found these regions activated in bvFTD as opposed to other FTD subtypes, possibly accounting for the effect on the pooled data.

We further found the strongest correlation of FTD and music perception in the insulae, which is in line with a previous report of involvement of the insula and putamen involved in hedonic musical changes in FTD (P. D. Fletcher et al. 2015). This activation was present in all FTD subtypes. Since the insula is one of the first brain areas that are affected in patients with FTD (Broe et al. 2003; Buhour et al. 2016), the reported changes in music perception could be an early indication of neurodegenerative processes in FTD, and possibly play a role in altered social cognitive abilities. Future research should further scrutinize the shared neurobiological mechanisms of music and social cognition in the disease pathogenesis of FTD.

A previous study by Seeley et al. observed that the brain regions atrophied in FTD are part of the functionally defined salience network, which has been shown to be disrupted in FTD patients and is important for complex social behavior (Seeley et al. 2009; Zhou et al. 2010). This network connects the insula and cingulate gyrus with the amygdala, thalamus and striatal regions. Our results show that atrophy patterns

Table 4

Results of the conjunction analyses. Illustrated are the clusters, the total volume of each cluster, x,y,z peak-coordinates and brain area. BA = Brodmann area.

Conjunction analysis of frontotemporal dementia and music perception					
Cluster	Volume (mm^3)	x	y	z	Label
1	9480	-32	14	10	Left insula BA 13
		-36	4	14	Left insula BA 13
		-36	24	10	Left inferior frontal gyrus BA 13
		-32	24	-4	Left inferior frontal gyrus BA 47
		-44	-2	0	Left insula BA 13
		-42	-10	4	Left insula BA 13
		-30	4	10	Left claustrum
		-42	30	4	Left inferior frontal gyrus BA 13
		-54	6	-10	Left superior temporal gyrus BA 38
		-46	12	4	Left insula BA 13
2	6552	32	32	-4	Right inferior frontal gyrus BA 47
		42	-4	-2	Right insula BA 13
		42	-10	2	Right insula BA 13
		36	22	8	Right inferior frontal gyrus BA 13
		44	-18	-2	Right insula BA 13
		34	26	-6	Right inferior frontal gyrus BA 47
		42	12	2	Right insula BA 13
		46	6	-4	Right insula BA 22
		46	12	-4	Right inferior frontal gyrus BA 47
		40	14	8	Right insula BA 13
3	2536	42	20	-6	Right inferior frontal gyrus BA 47
		34	12	16	Right insula BA 13
		16	4	-4	Right globus pallidus
		8	4	-4	Right caudate head
		8	6	2	Right caudate head
4	1512	10	6	6	Right caudate body
		-52	10	18	Left inferior frontal gyrus BA 44
		-50	12	24	Left inferior frontal gyrus BA 9
		-54	8	12	Left precentral gyrus BA 44
		-38	18	20	Left middle frontal gyrus BA 46
		-46	20	28	Left middle frontal gyrus BA 9
		-42	8	24	Left inferior frontal gyrus BA 9
5	1208	-52	-26	-2	Left superior temporal gyrus BA 21
		-46	-32	6	Left superior temporal gyrus BA 22
Conjunction analysis of frontotemporal dementia and social cognition					
Cluster	Volume (mm^3)	x	y	z	Label
1	27,432	-36	18	0	Left insula BA 13
		-22	-6	-14	Left amygdala
		-10	8	10	Left caudate body
		-40	2	0	Left insula BA 13
		-42	6	28	Left inferior frontal gyrus BA 9
		-46	8	-34	Left middle temporal gyrus BA 21
		-20	0	-6	Left lateral globus pallidus
		-20	4	6	Left putamen
		-48	12	-20	Left superior temporal gyrus BA 38
		-38	2	12	Left insula BA 13
		-32	32	-4	Left inferior frontal gyrus BA 47
		-54	8	8	Left precentral gyrus BA 44
		-38	-2	-12	Left superior temporal gyrus BA 21
		-46	0	38	Left middle frontal gyrus BA 6
		-52	10	18	Left inferior frontal gyrus BA 44

(continued on next page)

Table 4 (continued)

Conjunction analysis of frontotemporal dementia and music perception					
Cluster	Volume (mm ³)	x	y	z	Label
2	15,144	-4	50	8	Left medial frontal gyrus BA 10
		0	40	32	Left medial frontal gyrus BA 9
		0	38	36	Left medial frontal gyrus BA 6
		-6	16	40	Left cingulate gyrus BA 32
		-4	48	20	Left medial frontal gyrus BA 9
		0	26	44	Left medial frontal gyrus BA 8
		-4	30	24	Left anterior cingulate gyrus BA 32
		6	10	46	Right medial frontal gyrus BA 32
		-6	40	24	Left medial frontal gyrus BA 9
		-12	8	52	Left medial frontal gyrus BA 6
		34	18	-16	Right inferior frontal gyrus BA 47
		34	20	-8	Right inferior frontal gyrus BA 47
		36	18	-4	Right inferior frontal gyrus BA 47
		38	10	-32	Right superior temporal gyrus BA 38
3	10,552	40	-6	-4	Right insula BA 13
		44	20	-30	Right superior temporal gyrus BA 38
		48	-2	-22	Right fusiform gyrus BA 20
		48	8	-22	Right superior temporal gyrus BA 38
		34	30	-4	Right inferior frontal gyrus BA 47
		24	-8	-14	Right amygdala
		10	8	10	Right caudate body
		4	6	0	Right caudate head
		18	4	-4	Right putamen
		-54	-12	-22	Left inferior temporal gyrus BA 20
5	1736	-64	-10	-16	Left inferior temporal gyrus BA 21
		-52	-28	-4	Left middle temporal gyrus BA 21
6	1600	-52	-28	-4	Left middle temporal gyrus BA 21
7	736	0	-10	12	Left thalamus
8	488	0	38	-12	Left medial frontal gyrus BA 11
9	264	-40	16	48	Left superior frontal gyrus BA 8
10	256	46	-18	-8	Right superior temporal gyrus BA 22
Conjunction analysis of music perception and social cognition					
Cluster	Volume (mm ³)	x	y	z	Label
1	12,312	32	28	-2	Right inferior frontal gyrus BA 47
		52	4	-4	Right superior temporal gyrus BA 22
		44	-4	-4	Right insula BA 13
		46	-2	-8	Right superior temporal gyrus BA 22
		52	-24	2	Right superior temporal gyrus BA 22
		50	-22	-2	Right superior temporal gyrus BA 21
		42	20	30	Right middle frontal gyrus BA 9
		54	-10	-6	Right superior temporal gyrus BA 22
		56	-6	-6	Right middle temporal gyrus BA 21
		54	-2	-10	Right middle temporal gyrus BA 21
		52	-16	-4	Right superior temporal gyrus BA 22
		52	-28	-6	Right middle temporal gyrus BA 21
		60	-32	4	

Table 4 (continued)

Conjunction analysis of frontotemporal dementia and music perception					
Cluster	Volume (mm ³)	x	y	z	Label
2	11,200	48	26	8	Right middle temporal gyrus BA 22
		50	-2	-12	Right inferior frontal gyrus BA 45
		36	22	10	Right middle temporal gyrus BA 21
		42	12	4	Right insula BA 13
		58	20	22	Right inferior frontal gyrus BA 9
		42	22	-6	Right inferior frontal gyrus BA 47
		44	16	-6	Right inferior frontal gyrus BA 47
		-54	8	20	Left inferior frontal gyrus BA 44
		-32	14	12	Left insula BA 13
		-46	-2	-4	Left superior temporal gyrus BA 22
		-38	24	10	Left inferior frontal gyrus BA 13
		-34	0	12	Left insula BA 13
		-42	38	6	Left inferior frontal gyrus BA 46
		-32	24	-4	Left inferior frontal gyrus BA 47
		-46	30	8	Left inferior frontal gyrus BA 46
3	4144	-44	20	28	Left middle frontal gyrus BA 9
		-42	26	16	Left middle frontal gyrus BA 46
		-40	26	20	Left middle frontal gyrus BA 46
		-46	12	6	Left insula BA 13
		-42	8	24	Left inferior frontal gyrus BA 9
		-48	16	-6	Left inferior frontal gyrus BA 47
		-54	-26	2	Left superior temporal gyrus BA 22
		-54	-22	0	Left superior temporal gyrus BA 21
		-54	-12	-4	Left superior temporal gyrus BA 22
		-54	-6	-6	Left superior temporal gyrus BA 22
4	2216	-56	-42	20	Left superior temporal gyrus BA 13
		12	4	-4	Right globus pallidus
		10	6	2	Right caudate head
		10	-2	-4	Right medial globus pallidus
		50	-40	22	Right insula BA 13
5	1736	62	-40	22	Right superior temporal gyrus BA 22

observed in FTD mirror this network. Furthermore, we found that many of these regions are involved in music perception and social cognition. Dysfunction of this network is previously found to express with typical social-emotional problems (Farb et al. 2013; Rosen et al. 2002) just as music-emotion recognition (Omar et al. 2011) and musicophilia (Fletcher et al. 2013) in FTD patients. Finally, our subgroup analyses showed similar patterns of activation in FTD with both empathy and social cognition, particularly in the salience network. Our results provide further insight into social-emotional behavioral disturbances in FTD patients. These results have promising implications for the clinical applications of music as a probe for social cognitive and socio-emotional disturbances in diseases where these are most salient.

A potential limitation of our study is the translation of reported foci to voxels, which may have introduced noise into the results. The same principle applies to combining the data of structural and functional imaging studies. Another general limitation is heterogeneity of the

included studies. A minority of studies investigated genetic and anatomical variants of FTD. To improve statistical power we included studies with and without control conditions in the music perception group. Furthermore, as all the studies we included for music perception were listening tasks, and thus relatively homogenous in terms of tasks studied in contrast to social cognition. It is worth mentioning that the included studies predominantly used western music on western participants, and so it remains uncertain whether our conclusions are generalizable to non-western populations. A strength of our research was the large number of individuals that were included across all studies. Furthermore, we demonstrated an insightful and unique approach in investigating common neurobiological circuits by performing conjunction analyses on meta-analytic data.

5. Summary

Our meta-analysis demonstrates that music perception and social cognition share neurobiological circuits in frontotemporal dementia. This suggests music could be a sensitive probe for social cognition abilities with implications for diagnosis and monitoring.

CRediT authorship contribution statement

Jochum J. van't Hooft: Conceptualization, Investigation, Writing - original draft, Writing - review & editing, Visualization. **Yolande A.L. Pijnenburg:** Conceptualization, Writing - review & editing. **Sietske A.M. Sikkes:** Writing - review & editing. **Philip Scheltens:** Writing - review & editing. **Jacoba M. Spikman:** Writing - review & editing. **Artur C. Jaschke:** Writing - review & editing. **Jason D. Warren:** Writing - review & editing. **Betty M. Tijms:** Supervision, Conceptualization, Software, Writing - review & editing, Project administration, Resources.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. JW receives research grant support from the Alzheimer's Society and from the NIHR UCLH Biomedical Research Centre.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bandc.2020.105660>.

References

- Abell, F., Happé, F., & Frith, U. (2000). Do Triangles Play Tricks? Attribution of Mental States to Animated Shapes in Normal and Abnormal Development. *Cognitive Development*, 1, 1–22.
- Abraham, A., et al. (2008). Minds, Persons, and Space: An fMRI Investigation into the Relational Complexity of Higher-Order Intentionality. *Consciousness and Cognition*, 17(2), 438–450.
- Abraham, A., et al. (2010). Matching Mind to World and Vice Versa: Functional Dissociations between Belief and Desire Mental State Processing. *Social Neuroscience*, 5(1), 1–12.
- de Achával, D., et al. (2012). Decreased Activity in Right-Hemisphere Structures Involved in Social Cognition in Siblings Discordant for Schizophrenia. *Schizophrenia Research*, 141(1), 1402–1410.
- van Ackeren, M. J., Smaragdi, A., & Rueschemeyer, S. A. (2016). Neuronal Interactions between Mentalising and Action Systems during Indirect Request Processing. *Social Cognitive and Affective Neuroscience*, 11(9), 1402–1410.
- Adams, R. B., et al. (2010). Cross-Cultural Reading the Mind in the Eyes: An fMRI Investigation. *Journal of Cognitive Neuroscience*, 22(1), 97–108.
- Adenzato, M., Cavallo, M., & Enrici, I. (2010). Theory of Mind Ability in the Behavioural Variant of Frontotemporal Dementia: An Analysis of the Neural, Cognitive, and Social Levels. *Neuropsychologia*, 48(1), 2–12.
- Agosta, F., et al. (2012). White Matter Damage in Frontotemporal Lobar Degeneration Spectrum. *Cerebral Cortex*, 22(12), 2705–2714.
- Agosta, F., et al. (2009). Apolipoprotein E Epsilon4 Is Associated with Disease-Specific Effects on Brain Atrophy in Alzheimer's Disease and Frontotemporal Dementia. In *Proceedings of the National Academy of Sciences of the United States of America* (pp. 2018–2022).
- Agustus, J. L., et al. (2015). Functional MRI of Music Emotion Processing in Frontotemporal Dementia. *Annals of the New York Academy of Sciences*, 1337(1), 232–240.
- Ahmed, R. M., et al. (2016). "Assessment of Eating Behavior Disturbance and Associated Neural Networks in Frontotemporal Dementia". *JAMA Neurology*.
- Ahmed, R. M., et al. (2019). "Neural Networks Associated with Body Composition in Frontotemporal Dementia". *Annals of Clinical and Translational Neurology*.
- Aichhorn, M., et al. (2009). Temporo-Parietal Junction Activity in Theory-of-Mind Tasks: Falseness, Beliefs, or Attention. *Journal of Cognitive Neuroscience*, 21(6), 1179–1192.
- Aichhorn, M., et al. (2006). Do Visual Perspective Tasks Need Theory of Mind? *NeuroImage*, 30(3), 1059–1068.
- Akitsuki, Y., & Decety, J. (2009). Social Context and Perceived Agency Affects Empathy for Pain: An Event-Related fMRI Investigation. *NeuroImage*, 47(2), 722–734. <https://doi.org/10.1016/j.neuroimage.2009.04.091>.
- Alluri, V., et al. (2012). Large-Scale Brain Networks Emerge from Dynamic Processing of Musical Timbre, Key and Rhythm. *NeuroImage*, 59(4), 3677–3689.
- Altenmüller, E., et al. (2014). Play It Again, Sam: Brain Correlates of Emotional Music Recognition. *Frontiers in Psychology*, 5(FEB), 1–8.
- Angulo-Perkins, A., et al. (2014). Music Listening Engages Specific Cortical Regions within the Temporal Lobes: Differences between Musicians and Non-Musicians. *Cortex*, 50, 126–137.
- Aryani, A., Hsu, C. T., & Jacobs, A. M. (2019). Affective Iconic Words Benefit from Additional Sound-Meaning Integration in the Left Amygdala. *Human Brain Mapping*.
- Ash, S., et al. (2009). Non-Fluent Speech in Frontotemporal Lobar Degeneration. *Journal of Neurolinguistics*, 22(4), 370–383.
- Assaf, M., et al. (2009). Brain Activity Dissociates Mentalization from Motivation during an Interpersonal Competitive Game. *Brain Imaging and Behavior*, 3(1), 24–37.
- Azevedo, R. T., et al. (2014). Weighing the Stigma of Weight: An fMRI Study of Neural Reactivity to the Pain of Obese Individuals. *NeuroImage*, 91, 109–119. <https://doi.org/10.1016/j.neuroimage.2013.11.041>.
- Azevedo, R. T., et al. (2013). Their Pain Is Not Our Pain: Brain and Autonomic Correlates of Empathic Resonance with the Pain of Same and Different Race Individuals. *Human Brain Mapping*.
- Baez, S., Kanske, P., et al. (2016). Integration of Intention and Outcome for Moral Judgment in Frontotemporal Dementia: Brain Structural Signatures. *Neurodegenerative Diseases*, 16(3–4), 206–217.
- Baez, S., Morales, J. P., et al. (2016). Orbitofrontal and Limbic Signatures of Empathic Concern and Intentional Harm in the Behavioral Variant Frontotemporal Dementia. *Cortex*, 75, 20–32. <https://doi.org/10.1016/j.cortex.2015.11.007>.
- Baez, S., et al. (2019). Brain Structural Correlates of Executive and Social Cognition Profiles in Behavioral Variant Frontotemporal Dementia and Elderly Bipolar Disorder. *Neuropsychologia*, 126(January 2017), 159–169. <https://doi.org/10.1016/j.neuropsychologia.2017.02.012>.
- Bahnemann, M., et al. (2009). Sociotopy in the Temporoparietal Cortex: Common versus Distinct Processes. *Social Cognitive and Affective Neuroscience*, 5(1), 48–58.
- Bangert, M., et al. (2006). Shared Networks for Auditory and Motor Processing in Professional Pianists: Evidence from fMRI Conjunction. *NeuroImage*, 30(3), 917–926.
- Baron-Cohen, S., et al. (2000). The Amygdala Theory of Autism. *Neuroscience and Biobehavioral Reviews*, 24(3), 355–364.
- Baron-Cohen, S., et al. (2001). The 'Reading the Mind in the Eyes' Test Revised Version: A Study with Normal Adults, and Adults with Asperger Syndrome or High-Functioning Autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 42(2), 241–251.
- Baron-Cohen, S., Leslie, A. M., & Frith, U. (1985). Does the Autistic Child Have a 'Theory of Mind'? *Cognition*, 21(1), 37–46.
- Benuzzi, F., et al. (2018). Pain Mirrors: Neural Correlates of Observing Self or Others' Facial Expressions of Pain. *Frontiers in Psychology*, 9(OCT), 1–12.
- Benuzzi, F., et al. (2008). Does It Look Painful or Disgusting? Ask Your Parietal and Cingulate Cortex. *Journal of Neuroscience*, 28(4), 923–931.
- Berlingeri, M., et al. (2016). Guess Who's Coming to Dinner: Brain Signatures of Racially Biased and Politically Correct Behaviors. *Neuroscience*, 332, 231–241. <https://doi.org/10.1016/j.neuroscience.2016.06.048>.
- Bertoux, M., et al. (2018). Structural Anatomical Investigation of Long-Term Memory Deficit in Behavioral Frontotemporal Dementia. *Journal of Alzheimer's disease: JAD*, 62(4), 1887–1900.
- Berwick, R. C., Friederici, A. D., Chomsky, N., & Bolhuis, J. J. (2013). Evolution, Brain, and the Nature of Language. *Trends in Cognitive Sciences*, 17(2), 98.
- Bianco, R., et al. (2016). Neural Networks for Harmonic Structure in Music Perception and Action. *NeuroImage*, 142, 454–464.
- Boccadoro, S., et al. (2019). Defining the Neural Correlates of Spontaneous Theory of Mind (ToM): An fMRI Multi-Study Investigation. *NeuroImage*, 203(September), Article 116193. <https://doi.org/10.1016/j.neuroimage.2019.116193>.
- Boccardi, M., et al. (2005). Frontotemporal Dementia as a Neural System Disease. *Neurobiology of Aging*, 26(1), 37–44.

- Boden, M. E., et al. (2013). Comparing the Neural Correlates of Affective and Cognitive Theory of Mind Using fMRI: Involvement of the Basal Ganglia in Affective Theory of Mind. *Advances in Cognitive Psychology*, 9(1), 32–43.
- Boeve, B. F., & Geda, Y. E. (2001). Polka Music and Semantic Dementia. *Neurology*, 57, 1485.
- Bogert, B., et al. (2016). Hidden Sources of Joy, Fear, and Sadness: Explicit versus Implicit Neural Processing of Musical Emotions. *Neuropsychologia*, 89, 393–402. <https://doi.org/10.1016/j.neuropsychologia.2016.07.005>.
- Bora, E., Velakoulis, D., & Walterfang, M. (2016). Meta-Analysis of Facial Emotion Recognition in Behavioral Variant Frontotemporal Dementia. *Journal of Geriatric Psychiatry and Neurology*, 29(4), 205–211.
- Bos, P. A., et al. (2015). Oxytocin Reduces Neural Activity in the Pain Circuitry When Seeing Pain in Others. *NeuroImage*, 113, 217–224.
- Botvinick, M., et al. (2005). Viewing Facial Expressions of Pain Engages Cortical Areas Involved in the Direct Experience of Pain. *NeuroImage*, 25(1), 312–319.
- Boxer, A. L., et al. (2003). Cinguloparietal Atrophy Distinguishes Alzheimer Disease from Semantic Dementia. *Archives of Neurology*, 60(7), 949–956.
- Brambati, S. M., et al. (2009). Atrophy Progression in Semantic Dementia with Asymmetric Temporal Involvement: A Tensor-Based Morphometry Study. *Neurobiology of Aging*, 30(1), 103–111.
- Broe, M., et al. (2003). Staging Disease Severity in Pathologically Confirmed Cases of Frontotemporal Dementia. *Neurology*.
- Brown, S., & Martinez, M. J. (2007). Activation of Premotor Vocal Areas during Musical Discrimination. *Brain and Cognition*, 63(1), 59–69.
- Brüne, M., et al. (2008). An fMRI Study of Theory of Mind in Schizophrenic Patients with 'Passivity' Symptoms. *Neuropsychologia*, 46(7), 1992–2001.
- Brüne, M. (2005). Emotion Recognition, 'theory of Mind', and Social Behavior in Schizophrenia. *Psychiatry Research*.
- Bruneau, E. G., Jacoby, N., & Saxe, R. (2015). Empathic Control through Coordinated Interaction of Amygdala, Theory of Mind and Extended Pain Matrix Brain Regions. *NeuroImage*, 114, 105–119. <https://doi.org/10.1016/j.neuroimage.2015.04.034>.
- Bruneau, Emilie G., Agnieszka Pluta, and Rebecca Saxe. 2012. "Distinct Roles of the 'Shared Pain' and 'Theory of Mind' Networks in Processing Others' Emotional Suffering." *Neuropsychologia* 50(2): 219–31. <https://doi.org/10.1016/j.neuropsychologia.2011.11.008>.
- Brunet, E., Sarfati, Y., Hardy-Baylé, M. C., & Decety, J. (2000). A PET Investigation of the Attribution of Intentions with a Nonverbal Task. *NeuroImage*.
- Brünnlieb, C., Münte, T. F., Tempelmann, C., & Heldmann, M. (2013). Vasopressin Modulates Neural Responses Related to Emotional Stimuli in the Right Amygdala. *Brain Research*, 1499, 29–42.
- Budell, L., Jackson, P., & Rainville, P. (2010). Brain Responses to Facial Expressions of Pain: Emotional or Motor Mirroring? *NeuroImage*, 53(1), 355–363. <https://doi.org/10.1016/j.neuroimage.2010.05.037>.
- Buhour, M. S., et al. (2016). Pathophysiology of the Behavioral Variant of Frontotemporal Lobar Degeneration: A Study Combining MRI and FDG-PET. *Brain Imaging and Behavior*, 1–13.
- Buunk, A. M., et al. (2017). Social Cognition Impairments after Aneurysmal Subarachnoid Haemorrhage: Associations with Deficits in Interpersonal Behaviour, Apathy, and Impaired Self-Awareness. *Neuropsychologia*, 103(FEBRUARY), 131–139. <https://doi.org/10.1016/j.neuropsychologia.2017.07.015>.
- Cairns, Nigel J., et al. (2007). Neuropathologic diagnostic and nosologic criteria for frontotemporal lobar degeneration: consensus of the Consortium for Frontotemporal Lobar Degeneration. *Acta neuropathologica*, 114(5–22).
- Canessa, N., et al. (2012). The Neural Bases of Social Intention Understanding: The Role of Interaction Goals. *PLoS ONE*, 7(7).
- Cassidy, B., Hughes, C., & Krendl, A. (2020). Age Differences in Neural Activity Related to Mentalizing during Person Perception. *Aging, Neuropsychology, and Cognition*.
- Castelli, F., Happé, F., Frith, U., & Frith, C. (2000). Movement and Mind: A Functional Imaging Study of Perception and Interpretation of Complex Intentional Movement Patterns. *NeuroImage*.
- Castelli, I., et al. (2010). Effects of Aging on Mindreading Ability through the Eyes: An fMRI Study. *Neuropsychologia*, 48(9), 2586–2594. <https://doi.org/10.1016/j.neuropsychologia.2010.05.005>.
- Chaminade, T., et al. (2012). How Do We Think Machines Think? An fMRI Study of Alleged Competition with an Artificial Intelligence. *Frontiers in Human Neuroscience*, 6(MAY 2012), 1–9.
- Chapin, H., et al. (2010). Dynamic Emotional and Neural Responses to Music Depend on Performance Expression and Listener Experience. *PLoS ONE*, 5(12).
- Chen, J. L., Rae, C., & Watkins, K. E. (2012). Learning to Play a Melody: An fMRI Study Examining the Formation of Auditory-Motor Associations. *NeuroImage*, 59(2), 1200–1208.
- Cheng, Y., et al. (2010). Love Hurts: An fMRI Study. *NeuroImage*, 51(2), 923–929. <https://doi.org/10.1016/j.neuroimage.2010.02.047>.
- Cheng, Y., et al. (2007). Expertise Modulates the Perception of Pain in Others. *Current Biology*, 17(19), 1708–1713.
- Cheon, B. K., et al. (2013). Cultural Modulation of the Neural Correlates of Emotional Pain Perception: The Role of Other-Focusedness. *Neuropsychologia*, 51(7), 1177–1186. <https://doi.org/10.1016/j.neuropsychologia.2013.03.018>.
- Cheung, H., et al. (2012). False Belief and Verb Non-Factivity: A Common Neural Basis? *International Journal of Psychophysiology*, 83(3), 357–364. <https://doi.org/10.1016/j.ijpsycho.2011.12.002>.
- Chiao, Joao Y., Vani A. Mathur, Tokiko Harada, and Trixie Lipke. 2009. "Neural Basis of Preference for Human Social Hierarchy versus Egalitarianism." In Annals of the New York Academy of Sciences.
- Christov-Moore, L., & Iacoboni, M. (2019). Sex Differences in Somatomotor Representations of Others' Pain: A Permutation-Based Analysis. *Brain Structure and Function*.
- Clark, C. N., Downey, L. E., & Warren, J. D. (2014). Music Biology: All This Useful Beauty. *Current Biology*, 24(6).
- Clark, C. N., Downey, L. E., & Warren, J. D. (2015). Brain Disorders and the Biological Role of Music. *Social Cognitive and Affective Neuroscience*, 10(091673), 444–452. <http://www.ncbi.nlm.nih.gov/pubmed/24847111>.
- Contreras, J. M., Schirmer, J., Banaji, M. R., & Mitchell, J. P. (2013). Common Brain Regions with Distinct Patterns of Neural Responses during Mentalizing about Groups and Individuals. *Journal of Cognitive Neuroscience*.
- Corcoran, R., Mercer, G., & Frith, C. D. (1995). Schizophrenia, Symptomatology and Social Inference: Investigating 'Theory of Mind' in People with Schizophrenia. *Schizophrenia Research*, 17(1), 5–13.
- Corradi-Dell'acqua, Corrado, Hofstetter, Christoph, & Vuilleumier, Patrik (2011). Felt and Seen Pain Evoke the Same Local Patterns of Cortical Activity in Insular and Cingulate Cortex. *Journal of Neuroscience*, 31(49), 17996–18006.
- Corradi-D'Acqua, Corrado, Hofstetter, Christoph, & Vuilleumier, Patrik (2014). Cognitive and Affective Theory of Mind Share the Same Local Patterns of Activity in Posterior Temporal but Not Medial Prefrontal Cortex. *Social Cognitive and Affective Neuroscience*, 9(8), 1175–1184.
- Couto, Blas, et al. (2013). Structural Neuroimaging of Social Cognition in Progressive Non-Fluent Aphasia and Behavioral Variant of Frontotemporal Dementia. *Frontiers in Human Neuroscience*, 7.
- Danziger, Nicolas, Faillenot, Isabelle, & Peyron, Roland (2009). Can We Share a Pain We Never Felt? Neural Correlates of Empathy in Patients with Congenital Insensitivity to Pain. *Neuron*, 61(2), 203–212. <https://doi.org/10.1016/j.neuron.2008.11.023>.
- Deeley, Quinton, et al. (2006). Facial Emotion Processing in Criminal Psychopathy: Preliminary Functional Magnetic Resonance Imaging Study. *British Journal of Psychiatry*, 189(DEC), 533–539.
- Derntl, Birgit, et al. (2010). Multidimensional Assessment of Empathic Abilities: Neural Correlates and Gender Differences. *Psychoneuroendocrinology*, 35(1), 67–82.
- Desgranges, Béatrice, et al. (2007). Anatomical and Functional Alterations in Semantic Dementia: A Voxel-Based MRI and PET Study. *Neurobiology of Aging*, 28(12), 1904–1913.
- Deuse, L., et al. (2016). Neural Correlates of Naturalistic Social Cognition: Brain-Behavior Relationships in Healthy Adults. *Social Cognitive and Affective Neuroscience*, 11(11), 1741–1751.
- Dodell-Feder, David, Koster-Hale, Jorie, Bedny, Marina, & Saxe, Rebecca (2011). fMRI Item Analysis in a Theory of Mind Task. *NeuroImage*, 55(2), 705–712.
- Döhnel, K., et al. (2017). An fMRI Study on the Comparison of Different Types of False Belief Reasoning: False Belief-Based Emotion and Behavior Attribution. *Social Neuroscience*.
- Döhnel, Katrin, et al. (2012). Functional Activity of the Right Temporo-Parietal Junction and of the Medial Prefrontal Cortex Associated with True and False Belief Reasoning. *NeuroImage*, 60(3), 1652–1661.
- Downey, Laura E., et al. (2013). Mentalising Music in Frontotemporal Dementia. *Cortex*, 49(7), 1844–1855.
- Downey, Laura E., et al. (2015). White Matter Tract Signatures of Impaired Social Cognition in Frontotemporal Lobar Degeneration. *NeuroImage: Clinical*, 8, 640–651.
- Dufour, Nicholas, et al. (2013). Similar Brain Activation during False Belief Tasks in a Large Sample of Adults with and without Autism. *PLoS ONE*, 8(9).
- Dungan, James A., & Young, Liane (2019). Asking 'why?' Enhances Theory of Mind When Evaluating Harm but Not Purity Violations. *Social Cognitive and Affective Neuroscience*, 14(7), 699–708.
- Dziobek, Isabel, et al. (2006). Introducing MASC: A Movie for the Assessment of Social Cognition. *Journal of Autism and Developmental Disorders*.
- Eickhoff, S. B., et al. (2012). Activation Likelihood Estimation Meta-Analysis Revisited. *NeuroImage*, 59(3), 2349–2361. <https://doi.org/10.1016/j.neuroimage.2011.09.017>.
- Eickhoff, Simon B., et al. (2009). Coordinate-Based Activation Likelihood Estimation Meta-Analysis of Neuroimaging Data: A Random-Effects Approach Based on Empirical Estimates of Spatial Uncertainty. *Human Brain Mapping*, 30(9), 2907–2926.
- Enzi, Björn, Amirine, Scharbanu, & Brüne, Martin (2016). Empathy for Pain-Related Dorsolateral Prefrontal Activity Is Modulated by Angry Face Perception. *Experimental Brain Research*, 234(11), 3335–3345.
- Ernst, Jutta, et al. (2013). Interceptive Awareness Enhances Neural Activity during Empathy. *Human Brain Mapping*.
- Escoffier, Nicolas, Zhong, Jidan, Schirmer, Annett, & Qiu, Anqi (2013). Emotional Expressions in Voice and Music: Same Code, Same Effect? *Human Brain Mapping*, 34(8), 1796–1810.
- Eslinger, Paul J., et al. (2011). Correlates of Empathic Deficits in Frontotemporal Dementia. *Journal of Neuropsychiatry & Clinical Neurosciences*, 23(1), 74–82.
- Fan, Yang Teng, et al. (2014). Empathic Arousal and Social Understanding in Individuals with Autism: Evidence from fMRI and ERP Measurements. *Social Cognitive and Affective Neuroscience*, 9(8), 1203–1213.
- Farb, Norman A. S., et al. (2013). Abnormal Network Connectivity in Frontotemporal Dementia: Evidence for Prefrontal Isolation. *Cortex*, 49(7), 1856–1873.
- Feng, Chunliang, et al. (2016). Social Hierarchy Modulates Neural Responses of Empathy for Pain. *Social Cognitive and Affective Neuroscience*, 11(3), 485–495.
- Filippi, Massimo, et al. (2013). Functional Network Connectivity in the Behavioral Variant of Frontotemporal Dementia. *Cortex*, 49(9), 2389–2401.
- Flanagan, Emma C., et al. (2016). False Recognition in Behavioral Variant Frontotemporal Dementia and Alzheimer's Disease-Disinhibition or Amnesia? *Frontiers in Aging Neuroscience*, 8(JUN), 1–11.

- Fletcher, P. C., et al. (1995). Other Minds in the Brain: A Functional Imaging Study of 'Theory of Mind' in Story Comprehension. *Cognition*, 57(2), 109–128.
- Fletcher, Phillip D., et al. (2015). Auditory Hedonic Phenotypes in Dementia: Abbehavioural and Neuroanatomical Analysis. *Cortex*, 67, 95–105.
- Fletcher, Phillip D., Laura E. Downey, Pirada Witonpanich, and Jason D. Warren. 2013. "The Brain Basis of Musicophilia: Evidence from Frontotemporal Lobar Degeneration." *Frontiers in Psychology* 4. <http://journal.frontiersin.org/article/10.3389/fpsyg.2013.00347/abstract>.
- Flores-Gutiérrez, Enrique O., et al. (2007). Metabolic and Electric Brain Patterns during Pleasant and Unpleasant Emotions Induced by Music Masterpieces. *International Journal of Psychophysiology*, 65(1), 69–84.
- Fourie, Melike M., et al. (2017). Empathy and Moral Emotions in Post-Apartheid South Africa: An fMRI Investigation. *Social Cognitive and Affective Neuroscience*, 12(6), 881–892.
- Fox, Peter T., et al. (2005). BrainMap Taxonomy of Experimental Design: Description and Evaluation. *Human Brain Mapping*, 25(1), 185–198.
- Fox, Peter T., & Lancaster, Jack L. (2002). Mapping Context and Content: The BrainMap Model. *Nature Reviews Neuroscience*, 3(4), 319–321.
- Frank, Chiyo Kobayashi, Baron-Cohen, Simon, & Ganzel, Barbara L. (2015). Sex Differences in the Neural Basis of False-Belief and Pragmatic Language Comprehension. *NeuroImage*, 105, 300–311. <https://doi.org/10.1016/j.neuroimage.2014.09.041>.
- Fröhholz, Sascha, Trost, Wiebke, & Kotz, Sonja A. (2016). The Sound of Emotions—Towards a Unifying Neural Network Perspective of Affective Sound Processing. *Neuroscience and Biobehavioral Reviews*.
- Fujino, Junya, et al. (2014). Altered Brain Response to Others' Pain in Major Depressive Disorder. *Journal of Affective Disorders*, 165, 170–175. <https://doi.org/10.1016/j.jad.2014.04.058>.
- Gallagher, H., Happé, F., Brunswick, N., & Fletcher, P. (2000). Reading the Mind in Cartoons and Stories: An fMRI Study of 'theory of Mind' in Verbal. *Neuropsychologia*, 38, 11–21. [https://doi.org/10.1016/j.neuroimage.2019.116102](http://linkinghub.elsevier.com/retrieve/pii/S0028393299000536%5Cnfile:///Users/leeron/Dropbox/Papers2/Articles/Gallagher/2000/Gallagher_Reading_the_mind_in_cartoons_and_stories_an_fMRI_study_of_'theory_of_mind'_in_verbal_.?2000.pdf%5Cnpapers2:/publication.</p>
<p>Gallagher, Helen L., & Frith, Christopher D. (2004). Dissociable Neural Pathways for the Perception and Recognition of Expressive and Instrumental Gestures. <i>Neuropsychologia</i>, 42(13), 1725–1736.</p>
<p>García-Cordero, Indira, et al. (2015). Stroke and Neurodegeneration Induce Different Connectivity Aberrations in the Insula. <i>Stroke</i>, 46(9), 2673–2677.</p>
<p>Geiger, Alexander, et al. (2019). Distinct Functional Roles of the Mirror Neuron System and the Mentalizing System. <i>NeuroImage</i>, 202(July), Article 116102. <a href=).
- Geroldi, C., et al. (2000). Pop Music and Frontotemporal Dementia. *Neurology*, 55(12), 1935–1936.
- Gilbert, Sam J., et al. (2007). Distinct Regions of Medial Rostral Prefrontal Cortex Supporting Social and Nonsocial Functions. *Social Cognitive and Affective Neuroscience*, 2(3), 217–226.
- Gobbini, María Ida, et al. (2007). Two Takes on the Social Brain: A Comparison of Theory of Mind Tasks. *Journal of Cognitive Neuroscience*, 19(11), 1803–1814. <http://www.mitpressjournals.org/doi/10.1162/jocn.2007.19.11.1803>.
- Gorno-Tempini, M. L., et al. (2006). Anatomical Correlates of Early Mutism in Progressive Nonfluent Aphasia. *Neurology*.
- Gorno-Tempini, María Luisa, et al. 2004. Cognition and Anatomy in Three Variants of Primary Progressive Aphasia. *Annals of Neurology*.
- Gorno-Tempini, M. L., et al. (2011). Classification of primary progressive aphasia and its variants. *Neurology*, 76(11).
- Gossink, F., et al. (2018). Social Cognition Differentiates Behavioral Variant Frontotemporal Dementia From Other Neurodegenerative Diseases and Psychiatric Disorders. *American Journal of Geriatric Psychiatry*.
- Göttlich, Martin, et al. (2017). Viewing Socio-Affective Stimuli Increases Connectivity within an Extended Default Mode Network. *NeuroImage*, 148(December 2016), 8–19. <https://doi.org/10.1016/j.neuroimage.2016.12.044>.
- de Grecq, Moritz, et al. (2012). Neural Substrates Underlying Intentional Empathy. *Social Cognitive and Affective Neuroscience*, 7(2), 135–144.
- De Grecq, Moritz, et al. (2012). Altered Brain Activity during Emotional Empathy in Somatiform Disorder. *Human Brain Mapping*.
- Grézes, J., Pichon, S., & de Gelder, B. (2007). Perceiving Fear in Dynamic Body Expressions. *NeuroImage*, 35(2), 959–967.
- Grice-Jackson, Thomas, Critchley, Hugo D., Banissy, Michael J., & Ward, Jamie (2017). Consciously Feeling the Pain of Others Reflects Atypical Functional Connectivity between the Pain Matrix and Frontal-Parietal Regions. *Frontiers in Human Neuroscience*, 11(October), 1–13.
- Grossman, Murray, et al. (2004). What's in a Name: Voxel-Based Morphometric Analyses of MRI and Naming Difficulty in Alzheimer's Disease, Frontotemporal Dementia and Corticobasal Degeneration. *Brain*, 127(3), 628–649.
- Groussard, M., et al. (2010). The Neural Substrates of Musical Memory Revealed by fMRI and Two Semantic Tasks. *NeuroImage*, 53(4), 1301–1309.
- Groussard, Mathilde, et al. (2010). When Music and Long-Term Memory Interact: Effects of Musical Expertise on Functional and Structural Plasticity in the Hippocampus. *PLoS ONE*, 5(10).
- Gu, X., et al. (2013). Cognition-Emotion Integration in the Anterior Insular Cortex. *Cerebral Cortex*, 23(1), 20–27.
- Gu, Xiaosi, et al. (2010). Functional Dissociation of the Frontoinsular and Anterior Cingulate Cortices in Empathy for Pain. *Journal of Neuroscience*, 30(10), 3739–3744.
- Gu, Xiaosi, & Han, Shihui (2007). Attention and Reality Constraints on the Neural Processes of Empathy for Pain. *NeuroImage*, 36(1), 256–267. <https://doi.org/10.1016/j.neuroimage.2007.02.025>.
- Guevara, Andrea Brioschi, et al. (2015). Theory of Mind Impairment in Patients with Behavioural Variant Fronto-Temporal Dementia (Bv-FTD) Increases Caregiver Burden. *Age and Ageing*, 44(5), 891–895.
- Guo, Xiuyan, et al. (2013). Exposure to Violence Reduces Empathetic Responses to Other's Pain. *Brain and Cognition*, 82(2), 187–191. <https://doi.org/10.1016/j.bandc.2013.04.005>.
- Gweon, Hyowon, Dodell-Feder, David, Bedny, Marina, & Saxe, Rebecca (2012). Theory of Mind Performance in Children Correlates With Functional Specialization of a Brain Region for Thinking About Thoughts. *Child Development*, 83(6), 1853–1868.
- Haas, Brian W., Anderson, Ian W., & Filkowski, Megan M. (2015). Interpersonal Reactivity and the Attribution of Emotional Reactions. *Emotion*, 15(3), 390–398.
- Habermeyer, Benedikt, et al. (2009). Neural Correlates of Pre-Attentive Processing of Pattern Deviance in Professional Musicians. *Human Brain Mapping*, 30(11), 3736–3747.
- Hadjikhani, N. et al. 2014. "Emotional Contagion for Pain Is Intact in Autism Spectrum Disorders." *Translational Psychiatry* 4(November 2013).
- Han, Shihui, et al. (2009). Empathic Neural Responses to Others' Pain Are Modulated by Emotional Contexts. *Human Brain Mapping*.
- Han, Xiaochun, et al. (2017). Empathy for Pain Motivates Actions without Altruistic Effects: Evidence of Motor Dynamics and Brain Activity. *Social Cognitive and Affective Neuroscience*, 12(6), 893–901.
- Hardy, Chris J. D., et al. (2016). Hearing and Dementia. *Journal of Neurology*, 263(11), 2339–2354.
- Hardy, Chris JD, et al. (2017). Behavioural and Neuroanatomical Correlates of Auditory Speech Analysis in Primary Progressive Aphasias. *Alzheimer's Research and Therapy*, 9(1), 1–10.
- Harris, Lasana T., Todorov, Alexander, & Fiske, Susan T. (2005). Attributions on the Brain: Neuro-Imaging Dispositional Inferences, beyond Theory of Mind. *NeuroImage*, 28(4), 763–769.
- Harvey, R. J., Skelton-Robinson, M., & Rossor, M. N. (2003). The Prevalence and Causes of Dementia in People under the Age of 65 Years. *Journal of Neurology, Neurosurgery and Psychiatry*, 74(9), 1206–1209.
- Hervé, Pierre Yves, Razafimandimbify, Annick, Jobard, Gaël, & Tzourio-Mazoyer, Nathalie (2013). A Shared Neural Substrate for Mentalizing and the Affective Component of Sentence Comprehension. *PLoS ONE*, 8(1).
- Van Hoeck, Nicole, et al. (2014). False Belief and Counterfactual Reasoning in a Social Environment. *NeuroImage*, 90, 315–325. <https://doi.org/10.1016/j.neuroimage.2013.12.043>.
- Hooker, C., et al. (2010). Neural Activity during Social Signal Perception Correlates with Self-Reported Empathy. *Brain Research*, 1308, 100–113. <https://doi.org/10.1016/j.brainres.2009.10.006>.
- Hooker, Christine I., et al. (2008). Mentalizing about Emotion and Its Relationship to Empathy. *Social Cognitive and Affective Neuroscience*, 3(3), 204–217.
- Hornberger, Michael, Geng, John, & Hodges, John R. (2011). Convergent Grey and White Matter Evidence of Orbitofrontal Cortex Changes Related to Disinhibition in Behavioural Variant Frontotemporal Dementia. *Brain*, 134(9), 2502–2512.
- Hsieh, Sharpley, et al. (2013). When One Loses Empathy: Its Effect on Carers of Patients with Dementia. *Journal of Geriatric Psychiatry and Neurology*, 26(3), 174–184.
- Irish, Muireann, Eyre, Nadine, et al. (2016). Neural Substrates of Semantic Prospection – Evidence from the Dementias. *Frontiers in Behavioral Neuroscience*, 10(MAY), 1–12.
- Irish, Muireann, Bunk, Steffie, et al. (2016). Preservation of Episodic Memory in Semantic Dementia: The Importance of Regions beyond the Medial Temporal Lobes. *Neuropsychologia*, 81, 50–60.
- Jackson, Philip L., Brunet, Eric, Meltzoff, Andrew N., & Decety, Jean (2006). Empathy Examined through the Neural Mechanisms Involved in Imagining How I Feel versus How You Feel Pain. *Neuropsychologia*, 44(5), 752–761.
- Jackson, Philip L., Meltzoff, Andrew N., & Decety, Jean (2005). How Do We Perceive the Pain of Others? A Window into the Neural Processes Involved in Empathy. *NeuroImage*, 24(3), 771–779.
- Jacoby, Nir, Bruneau, Emile, Koster-Hale, Jorie, & Saxe, Rebecca (2016). Localizing Pain Matrix and Theory of Mind Networks with Both Verbal and Non-Verbal Stimuli. *NeuroImage*.
- Janata, P. (2015). Neural Basis of Music Perception. *Handbook of Clinical Neurology*, 129, 187–205.
- Janata, Petr, Birk, Jeffrey L., et al. (2002). The Cortical Topography of Tonal Structures Underlying Western Music. *Science*, 298(5601), 2167–2170.
- Janata, Petr (2009). The Neural Architecture of Music-Evoked Autobiographical Memories. *Cerebral Cortex*, 19(11), 2579–2594.
- Janata, Petr, Tillmann, Barbara, & Bharucha, Jamshed J. (2002). Listening to Polyphonic Music Recruits Domain-General Attention and Working Memory Circuits. *Cognitive, Affective and Behavioral Neuroscience*, 2(2), 121–140.
- Jankowiak-Siuda, Kamila et al. 2015. Physical Attractiveness and Sex as Modulatory Factors of Empathic Brain Responses to Pain. *Frontiers in Behavioral Neuroscience*.
- Jenkins, Adrianna C., & Mitchell, Jason P. (2010). Mentalizing under Uncertainty: Dissociated Neural Responses to Ambiguous and Unambiguous Mental State Inferences. *Cerebral Cortex*, 20(2), 404–410.
- Jimura, Koji, Konishi, Seiki, Asari, Tomoki, & Miyashita, Yasushi (2010). Temporal Pole Activity during Understanding Other Persons' Mental States Correlates with Neuroticism Trait. *Brain Research*, 1328, 104–112. <https://doi.org/10.1016/j.brainres.2010.03.016>.
- Kaiser, Stefan, et al. (2008). Gender-Specific Strategy Use and Neural Correlates in a Spatial Perspective Taking Task. *Neuropsychologia*, 46(10), 2524–2531.

- Kana, Rajesh, et al. (2016). Altered Medial Frontal and Superior Temporal Response to Implicit Processing of Emotions in Autism. *Autism Research*.
- Kana, Rajesh K., et al. (2009). Atypical Frontal-Posterior Synchronization of Theory of Mind Regions in Autism during Mental State Attribution. *Social Neuroscience*.
- Kana, Rajesh K., & Travers, Brittany G. (2012). Neural Substrates of Interpreting Actions and Emotions from Body Postures. *Social Cognitive and Affective Neuroscience*, 7(4), 446–456.
- Kanda, Tomonori, et al. (2008). Comparison of Grey Matter and Metabolic Reductions in Frontotemporal Dementia Using FDG-PET and Voxel-Based Morphometric MR Studies. *European Journal of Nuclear Medicine and Molecular Imaging*, 35(12), 2227–2234.
- Kandylaki, Katerina D., et al. (2015). Processing of False Belief Passages during Natural Story Comprehension: An fMRI Study. *Human Brain Mapping*, 36(11), 4231–4246.
- Kanske, Philipp, Böckler, Anne, Trautwein, Fynn Mathis, & Singer, Tania (2015). Dissecting the Social Brain: Introducing the EmpaToM to Reveal Distinct Neural Networks and Brain-Behavior Relations for Empathy and Theory of Mind. *NeuroImage*, 122, 6–19. <https://doi.org/10.1016/j.neuroimage.2015.07.082>.
- Kim, E. J., et al. (2007). Patterns of MRI Atrophy in Tau Positive and Ubiquitin Positive Frontotemporal Lobar Degeneration. *Journal of Neurology, Neurosurgery and Psychiatry*, 78(12), 1375–1378.
- Kim, Eosu, et al. (2009). Reduced Activation in the Mirror Neuron System during a Virtual Social Cognition Task in Euthymic Bipolar Disorder. *Progress in Neuropsychopharmacology and Biological Psychiatry*, 33(8), 1409–1416. <https://doi.org/10.1016/j.pnpbp.2009.07.019>.
- Kim, Ji Woong, et al. (2005). Neural Mechanism for Judging the Appropriateness of Facial Affect. *Cognitive Brain Research*, 25(3), 659–667.
- Kim, Sung Eun, et al. (2007). The Neural Mechanism of Imagining Facial Affective Expression. *Brain Research*, 1145(1), 128–137.
- Kipps, C. M., et al. (2009). Understanding Social Dysfunctional in the Behavioural Variant of Frontotemporal Dementia: The Role of Emotion and Sarcasm Processing. *Brain*, 132(3), 592–603.
- Kircher, Tilo, et al. (2009). Online Mentalising Investigated with Functional MRI. *Neuroscience Letters*, 454(3), 176–181.
- Klimecki, Olga M., Leiberg, Susanne, Ricard, Matthieu, & Singer, Tania (2013). Differential Pattern of Functional Brain Plasticity after Compassion and Empathy Training. *Social Cognitive and Affective Neuroscience*.
- Kobayashi, Chiyo, Glover, Gary H., & Temple, Elise (2008). Switching Language Switches Mind: Linguistic Effects on Developmental Neural Bases of 'Theory of Mind'. *Social Cognitive and Affective Neuroscience*, 3(1), 62–70.
- Kockler, H., et al. (2010). Visuospatial Perspective Taking in a Dynamic Environment: Perceiving Moving Objects from a First-Person-Perspective Induces a Disposition to Act. *Consciousness and Cognition*, 19(3), 690–701. <https://doi.org/10.1016/j.concog.2010.03.003>.
- Koelkebeck, Katja, et al. (2011). Transcultural Differences in Brain Activation Patterns during Theory of Mind (ToM) Task Performance in Japanese and Caucasian Participants. *Social Neuroscience*.
- Krach, et al. (2015). Evidence from Pupilometry and fMRI Indicates Reduced Neural Response during Vicarious Social Pain but Not Physical Pain in Autism. *Human Brain Mapping*.
- Krach, S., et al. (2009). Are Women Better Mindreaders? Sex Differences in Neural Correlates of Mentalizing Detected with Functional MRI. *BMC Neuroscience*, 10, 1–11.
- Krach, S., et al. (2011). Your Flaws Are My Pain: Linking Empathy to Vicarious Embarrassment. *PLoS ONE*.
- Krach, Sören, et al. (2008). Can Machines Think? Interaction and Perspective Taking with Robots Investigated via fMRI. *PLoS ONE*, 3(7).
- Krämer, Ulrike M., et al. (2010). Emotional and Cognitive Aspects of Empathy and Their Relation to Social Cognition - an fMRI-Study. *Brain Research*, 1311, 110–120.
- Kumfor, Fiona, et al. (2018). Beyond the Face: How Context Modulates Emotion Processing in Frontotemporal Dementia Subtypes. *Brain*, 141(4), 1172–1185.
- Lagarde, J., et al. (2015). Why Do Patients with Neurodegenerative Frontal Syndrome Fail to Answer: 'In What Way Are an Orange and a Banana Alike?'. *Brain*, 138(2), 456–471.
- Lagarde, Julien, et al. (2013). Are Frontal Cognitive and Atrophy Patterns Different in PSP and BvFTD? A Comparative Neuropsychological and VBM Study. *PLoS ONE*, 8 (11).
- Laird, Angela R., Lancaster, Jack L., & Fox, Peter T. (2005). BrainMap: The Social Evolution of a Human Brain Mapping Database. *Neuroinformatics*, 3(1), 065–078. <http://link.springer.com/10.1385/NI:3:1:065>.
- Lamm, Claus, & Decety, Jean (2008). Is the Extrastriate Body Area (EBA) Sensitive to the Perception of Pain in Others? *Cerebral Cortex*, 18(10), 2369–2373.
- Lancaster, Jack L., et al. (2007). Bias between MNI and Talairach Coordinates Analyzed Using the ICBM-152 Brain Template. *Human Brain Mapping*, 28(11), 1194–1205.
- Lancaster, Katie, et al. (2015). Plasma Oxytocin Explains Individual Differences in Neural Substrates of Social Perception. *Frontiers in Human Neuroscience*, 9(MAR), 1–7.
- Langdon, R., Coltheart, M., Ward, P. B., & Catts, S. V. (2002). Disturbed Communication in Schizophrenia: The Role of Poor Pragmatics and Poor Mind-Reading. *Psychological Medicine*, 32(7), 1273–1284.
- Langheim, Frederick J. P., et al. (2002). Cortical Systems Associated with Covert Music Rehearsals. *NeuroImage*, 16(4), 901–908.
- Lavoie, Marie Audrey, et al. (2016). Social Representations and Contextual Adjustments as Two Distinct Components of the Theory of Mind Brain Network: Evidence from the REMICS Task. *Cortex*, 81, 176–191.
- Lawrence, E. J., et al. (2006). The Role of 'shared Representations' in Social Perception and Empathy: An fMRI Study. *NeuroImage*, 29(4), 1173–1184.
- Lee, Seung Jae, et al. (2010). Multi-Level Comparison of Empathy in Schizophrenia: An fMRI Study of a Cartoon Task. *Psychiatry Research - Neuroimaging*, 181(2), 121–129. <https://doi.org/10.1016/j.pscychresns.2009.08.003>.
- Lee, Suzee E., et al. (2014). Altered Network Connectivity in Frontotemporal Dementia with C9orf72 Hexanucleotide Repeat Expansion. *Brain*, 137(11), 3047–3060.
- Lee, Tatia M. C., et al. (2013). Neural Activities during Affective Processing in People with Alzheimer's Disease. *Neurobiology of Aging*, 34(3), 706–715.
- Leiberg, Susanne, Eippert, Falk, Veit, Ralf, & Anders, Silke (2012). Intentional Social Distance Regulation Alters Affective Responses towards Victims of Violence: An fMRI Study. *Human Brain Mapping*.
- Leitman, David I., et al. (2010). "It's Not What You Say, but How You Say It": A Reciprocal Temporo-Frontal Network for Affective Prosody. *Frontiers in Human Neuroscience*, 4(February), 1–13.
- Levitin, Daniel J., & Menon, Vinod (2003). Musical Structure Is Processed in 'Language' Areas of the Brain: A Possible Role for Brodmann Area 47 in Temporal Coherence. *NeuroImage*, 20(4), 2142–2152.
- Lewis, Penelope A., Birch, Amy, Hall, Alexander, & Dunbar, Robin I. M. (2017). Higher Order Intentionality Tasks Are Cognitively More Demanding. *Social Cognitive and Affective Neuroscience*, 12(7), 1063–1071.
- Libon, D. J., et al. (2009). Neurocognitive Contributions to Verbal Fluency Deficits in Frontotemporal Lobar Degeneration. *Neurology*, 73(7), 535–542.
- Liew, Sook Lei, Han, Shihui, & Aziz-Zadeh, Lisa (2011). Familiarity Modulates Mirror Neuron and Mentalizing Regions during Intention Understanding. *Human Brain Mapping*.
- Limb, Charles J. (2006). Structural and Functional Neural Correlates of Music Perception. *Anatomical Record - Part A Discoveries in Molecular, Cellular, and Evolutionary Biology*, 288(4), 435–446.
- Lin, Nan, et al. (2018). Neural Correlates of Three Cognitive Processes Involved in Theory of Mind and Discourse Comprehension. *Cognitive, Affective and Behavioral Neuroscience*, 18(2), 273–283.
- Lombardo, Michael V., et al. (2010). Shared Neural Circuits for Mentalizing about the Self and Others. *Journal of Cognitive Neuroscience*, 22(7), 1623–1635.
- Lotze, M., et al. (2006). Differential Cerebral Activation during Observation of Expressive Gestures and Motor Acts. *Neuropsychologia*, 44(10), 1787–1795.
- Luo, Chunyan, et al. (2020). Comparison of Gray Matter Atrophy in Behavioral Variant Frontal Temporal Dementia and Amyotrophic Lateral Sclerosis: A Coordinate-Based Meta-Analysis. *Frontiers in Aging Neuroscience*, 12(February).
- Luo, Siyang, et al. (2014). Reminders of Mortality Decrease Midcingulate Activity in Response to Others' Suffering. *Social Cognitive and Affective Neuroscience*, 9(4), 477–486.
- Malhi, Gin S., et al. (2008). A Functional MRI Study of Theory of Mind in Euthymic Bipolar Disorder Patients. *Bipolar Disorders*.
- Mandelli, Maria Luisa, et al. (2016). Two Insular Regions Are Differentially Involved in Behavioral Variant FTD and Nonfluent/Agrammatic Variant PPA. *Cortex*.
- Mano, Yoko, et al. (2009). Perspective-Taking as Part of Narrative Comprehension: A Functional MRI Study. *Neuropsychologia*, 47(3), 813–824.
- Marjoram, Dominic, et al. (2006). A Visual Joke fMRI Investigation into Theory of Mind and Enhanced Risk of Schizophrenia. *NeuroImage*, 31(4), 1850–1858.
- Martin, Alex, & Weisberg, Jill (2003). Neural Foundations for Understanding Social and Mechanical Concepts. *Cognitive Neuropsychology*.
- Mathur, Vani A., et al. (2016). Overlapping Neural Response to the Pain or Harm of People, Animals, and Nature. *Neuropsychologia*, 81, 265–273. <https://doi.org/10.1016/j.neuropsychologia.2015.12.025>.
- Mathur, Vani A., Harada, Tokiko, Lipke, Trixie, & Chiao, Joan Y. (2010). Neural Basis of Extraordinary Empathy and Altruistic Motivation. *NeuroImage*, 51(4), 1468–1475. <https://doi.org/10.1016/j.neuroimage.2010.03.025>.
- May, Michelle, et al. (2017). Social Behavior and Impairments in Social Cognition Following Traumatic Brain Injury. *Journal of the International Neuropsychological Society*, 23(5), 400–411.
- Mazza, Monica, et al. (2013). Regional Cerebral Changes and Functional Connectivity during the Observation of Negative Emotional Stimuli in Subjects with Post-Traumatic Stress Disorder. *European Archives of Psychiatry and Clinical Neuroscience*, 263(7), 575–583.
- Mazzola, Viridiana, et al. (2010). Affective Response to a Loved One's Pain: Insula Activity as a Function of Individual Differences. *PLoS ONE*, 5(12).
- McAdams, Carrie J., & Krawczyk, Daniel C. (2011). Impaired Neural Processing of Social Attribution in Anorexia Nervosa. *Psychiatry Research - Neuroimaging*, 194(1), 54–63. <https://doi.org/10.1016/j.pscychresns.2011.06.016>.
- McKhann, Guy M., et al. (2001). Clinical and pathological diagnosis of frontotemporal dementia: Report of the Work Group on Frontotemporal Dementia and Pick's Disease. *Archives of Neurology*, 58(1803–9).
- Meaux, Emilie, & Vuilleumier, Patrik (2016). Facing Mixed Emotions: Analytic and Holistic Perception of Facial Emotion Expressions Engages Separate Brain Networks. *NeuroImage*, 141, 154–173. <https://doi.org/10.1016/j.neuroimage.2016.07.004>.
- Van der Meer, Lisette, et al. (2011). Inhibit Yourself and Understand the Other: Neural Basis of Distinct Processes Underlying Theory of Mind. *NeuroImage*, 56(4), 2364–2374.
- Melchers, Martin, et al. (2015). Reality TV and Vicarious Embarrassment: An fMRI Study. *NeuroImage*, 109, 109–117. <https://doi.org/10.1016/j.neuroimage.2015.01.022>.
- Melloni, Margherita, et al. (2016). Your Perspective and My Benefit: Multiple Lesion Models of Self-Other Integration Strategies during Social Bargaining. *Brain*, 139(11), 3022–3040.
- Mercadillo, Roberto E., Díaz, José Luis, Pasaye, Erick H., & Barrios, Fernando A. (2011). Perception of Suffering and Compassion Experience: Brain Gender Disparities. *Brain and Cognition*, 76(1), 5–14. <https://doi.org/10.1016/j.bandc.2011.03.019>.

- Meyer, Meghan L., Taylor, Shelley E., & Lieberman, Matthew D. (2015). Social Working Memory and Its Distinctive Link to Social Cognitive Ability: An fMRI Study. *Social Cognitive and Affective Neuroscience*, 10(10), 1338–1347.
- Mier, D., et al. (2010). Neuronal Correlates of Affective Theory of Mind in Schizophrenia Out-Patients: Evidence for a Baseline Deficit. *Psychological Medicine*, 40(10), 1607–1617.
- Mier, Daniela, et al. (2010). The Involvement of Emotion Recognition in Affective Theory of Mind. *Psychophysiology*, 47(6), 1028–1039.
- Mitchell, Jason P. (2008). Activity in Right Temporo-Parietal Junction Is Not Selective for Theory-of-Mind. *Cerebral Cortex*, 18(2), 262–271.
- Mitterschiffthaler, Martina T., et al. (2007). A Functional MRI Study of Happy and Sad Affective States Induced by Classical Music. *Human Brain Mapping*, 28(11), 1150–1162.
- Moessnang, Carolin, et al. (2017). Differential Responses of the Dorsomedial Prefrontal Cortex and Right Posterior Superior Temporal Sulcus to Spontaneous Mentalizing. *Human Brain Mapping*.
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *BMJ*, 339(jul21 1), b2535–b2535.
- Molenberghs, Pascal, Johnson, Halle, Henry, Julie D., & Mattingley, Jason B (2016). Neuroscience and Biobehavioral Reviews Understanding the Minds of Others: A Neuroimaging Meta-Analysis. *Neuroscience and Biobehavioral Reviews*, 65, 276–291. <https://doi.org/10.1016/j.neubiorev.2016.03.020>.
- Moll, Jorge, et al. (2007). The Self as a Moral Agent: Linking the Neural Bases of Social Agency and Moral Sensitivity. *Social Neuroscience*.
- Moran, Joseph M., Jolly, Eshin, & Mitchell, Jason P. (2012). Social-Cognitive Deficits in Normal Aging. *Journal of Neuroscience*, 32(16), 5553–5561.
- Morelli, Sylvia A., Rameson, Lian T., & Lieberman, Matthew D. (2014). The Neural Components of Empathy: Predicting Daily Prosocial Behavior. *Social Cognitive and Affective Neuroscience*, 9(1), 39–47.
- Moriguchi, Yoshiya, et al. (2007). Empathy and Judging Other's Pain: An fMRI Study of Alexithymia. *Cerebral Cortex*, 17(9), 2223–2234.
- Morrison, India, & Downing, Paul E. (2007). Organization of Felt and Seen Pain Responses in Anterior Cingulate Cortex. *NeuroImage*, 37(2), 642–651.
- Morrison, India, Lloyd, Donna, Di Pellegrino, Giuseppe, & Roberts, Neil (2004). Vicarious Responses to Pain in Anterior Cingulate Cortex: Is Empathy a Multisensory Issue? *Cognitive, Affective and Behavioral Neuroscience*, 4(2), 270–278.
- Morrison, India, Peelen, Marius V., & Downing, Paul E. (2007). The Sight of Others' Pain Modulates Motor Processing in Human Cingulate Cortex. *Cerebral Cortex*, 17(9), 2214–2222.
- Morrison, India, Tipper, Steve P., Fenton-Adams, Wendy L., & Bach, Patric (2013). 'Feeling' Others' Painful Actions: The Sensorimotor Integration of Pain and Action Information. *Human Brain Mapping*.
- Morrison, Steven J., et al. (2003). fMRI Investigation of Cross-Cultural Music Comprehension. *NeuroImage*, 20(1), 378–384.
- Mummery, C. J., et al. (2000). A Voxel-Based Morphometry Study of Semantic Dementia: Relationship between Temporal Lobe Atrophy and Semantic Memory. *Annals of Neurology*, 47(1), 36–45.
- Murphy, Eric R., et al. (2010). Differential Processing of Metacognitive Evaluation and the Neural Circuitry of the Self and Others in Schizophrenia: A Pilot Study. *Schizophrenia Research*, 116(2–3), 252–258. <https://doi.org/10.1016/j.schres.2009.11.009>.
- Musso, Mariacristina, et al. (2015). A Single Dual-Stream Framework for Syntactic Computations in Music and Language. *NeuroImage*, 117, 267–283.
- Narumoto, Jin, et al. (2000). Brain Regions Involved in Verbal or Non-Verbal Aspects of Facial Emotion Recognition. *NeuroReport*, 11(11), 2571–2576.
- Neary, D., et al. (1998). Frontotemporal lobar degeneration: A consensus on clinical diagnostic criteria. *Neurology*, 51(6).
- Nijssse, Britta, et al. (2019). Social Cognition Impairments in the Long Term Post Stroke. *Archives of Physical Medicine and Rehabilitation*, 100(7), 1300–1307. <https://doi.org/10.1016/j.apmr.2019.01.023>.
- Normi, Jason S., et al. (2008). On the Neural Networks of Empathy: A Principal Component Analysis of an fMRI Study. *Behavioral and Brain Functions*, 4, 1–13.
- Nummenmaa, Lauri, Hirvonen, Jussi, Parkkola, Riitta, & Hietanen, Jari K. (2008). Is Emotional Contagion Special? An fMRI Study on Neural Systems for Affective and Cognitive Empathy. *NeuroImage*, 43(3), 571–580. <https://doi.org/10.1016/j.neuroimage.2008.08.014>.
- Ochsner, Kevin N., et al. (2004). Reflecting upon Feelings: An fMRI Study of Neural Systems Supporting the Attribution of Emotion to Self and Other. *Journal of Cognitive Neuroscience*, 16(10), 1746–1772. <http://www.mitpressjournals.org/doi/10.1162/0898929042947829>.
- Oechslin, Mathias S., Gschwind, Markus, & James, Clara E. (2017). Tracking Training-Related Plasticity by Combining fMRI and DTI: The Right Hemisphere Ventral Stream Mediates Musical Syntax Processing. *Cerebral Cortex*, 1–10. <https://academic.oup.com/cercor/article/2998920/Tracking>.
- Omar, Rohani, et al. (2011). The Structural Neuroanatomy of Music Emotion Recognition: Evidence from Frontotemporal Lobar Degeneration. *NeuroImage*, 56(3), 1814–1821. <https://doi.org/10.1016/j.neuroimage.2011.03.002>.
- Otti, Alexander, Wohlschlaeger, Afra M., & Noll-Hussong, Michael (2015). Is the Medial Prefrontal Cortex Necessary for Theory of Mind? *PLoS ONE*, 10(8).
- Van Overwalle, Frank, Daes, Tine, & Mariën, Peter (2015). Social Cognition and the Cerebellum: A Meta-Analytic Connectivity Analysis. *Human Brain Mapping*.
- Özdem, Ceylan, Brass, Marcel, Van der Cruyssen, Laurens, & Van Overwalle, Frank (2017). The Overlap between False Belief and Spatial Reorientation in the Temporo-Parietal Junction: The Role of Input Modality and Task. *Social Neuroscience*.
- Pan, Ping Lei, et al. (2012). Gray Matter Atrophy in Behavioral Variant Frontotemporal Dementia: A Meta-Analysis of Voxel-Based Morphometry Studies. *Dementia and Geriatric Cognitive Disorders*, 33(2–3), 141–148.
- Pardini, Matteo, Huey, Edward D., Cavanagh, Alyson L., & Grafman, Jordan (2009). Olfactory Function in Corticobasal Syndrome and Frontotemporal Dementia. *Archives of Neurology*, 66(1), 92–96.
- Park, Mona, et al. (2014). Differences between Musicians and Non-Musicians in Neuro-Affective Processing of Sadness and Fear Expressed in Music. *Neuroscience Letters*, 566, 120–124.
- Paulus, Frieder M., et al. (2018). Laugh or Cringe? Common and Distinct Processes of Reward-Based Schadenfreude and Empathy-Based Fremdscham. *Neuropsychologia*, 116(May 2017), 52–60. <https://doi.org/10.1016/j.neuropsychologia.2017.05.030>.
- Pereira, J. M. S., et al. (2009). Atrophy Patterns in Histologic vs Clinical Groupings of Frontotemporal Lobar Degeneration. *Neurology*, 72(19), 1653–1660.
- Pichon, Swann, de Gelder, Beatrice, & Grézes, Julie (2008). Emotional Modulation of Visual and Motor Areas by Dynamic Body Expressions of Anger. *Social Neuroscience*.
- Piguet, Olivier, Hornberger, Michael, Mioshi, Eneida, & Hodges, John R. (2011). Behavioural-Variant Frontotemporal Dementia: Diagnosis, Clinical Staging, and Management. *The Lancet Neurology*, 10(2), 162–172.
- Platek, Steven M., Keenan, Julian Paul, Gallup, Gordon G., & Mohamed, Feroze B. (2004). Where Am I? The Neurological Correlates of Self and Other. *Cognitive Brain Research*, 19(2), 114–122.
- Powell, Joanne L., et al. (2017). The Neural Correlates of Theory of Mind and Their Role during Empathy and the Game of Chess: A Functional Magnetic Resonance Imaging Study. *Neuroscience*, 355, 149–160.
- Preis, Mira A., Schmidt-Samoa, Carsten, Dechant, Peter, & Kroener-Herwig, Birgit (2013). The Effects of Prior Pain Experience on Neural Correlates of Empathy for Pain: An FMRI Study. *Pain*.
- Prochnow, D., et al. (2013). The Neural Correlates of Affect Reading: An fMRI Study on Faces and Gestures. *Behavioural Brain Research*, 237(1), 270–277. <https://doi.org/10.1016/j.bbr.2012.08.050>.
- Prochnow, D., Brunheim, S., Steinhäuser, L., & Seitz, R. J. (2014). Reasoning about the Implications of Facial Expressions: A Behavioral and fMRI Study on Low and High Social Impact. *Brain and Cognition*, 90, 165–173. <https://doi.org/10.1016/j.bandc.2014.07.004>.
- Qiao-Tassert, Emilie, Corradi-Dell'Acqua, Corrado, & Vuilleumier, Patrik (2018). The Good, the Bad, and the Suffering: Transient Emotional Episodes Modulate the Neural Circuits of Pain and Empathy. *Neuropsychologia*, 116(December 2017), 99–116. <https://doi.org/10.1016/j.neuropsychologia.2017.12.027>.
- Rabin, Jennifer S., et al. (2010). Common and Unique Neural Correlates of Autobiographical Memory and Theory of Mind. *Journal of Cognitive Neuroscience*, 22 (6), 1095–1111.
- Rabinovici, G. D., et al. (2008). Distinct MRI Atrophy Patterns in Autopsy-Proven Alzheimer's Disease and Frontotemporal Lobar Degeneration. *American Journal of Alzheimer's Disease and other Dementias*, 22(6), 474–488.
- Ranasinghe, Kamalini G., et al. (2016). Distinct Subtypes of Behavioral Variant Frontotemporal Dementia Based on Patterns of Network Degeneration. *JAMA Neurology*, 73(9), 1078–1088.
- Rankin, Katherine P., et al. (2011). Behavioral Variant Frontotemporal Dementia with Corticobasal Degeneration Pathology: Phenotypic Comparison to BvFTD with Pick's Disease. *Journal of Molecular Neuroscience*.
- Rapp, A. M., et al. (2010). Neural Correlates of Irony Comprehension: The Role of Schizotypal Personality Traits. *Brain and Language*, 113(1), 1–12. <https://doi.org/10.1016/j.bandl.2009.11.007>.
- Rascovsky, Katya, et al. (2011). Sensitivity of Revised Diagnostic Criteria for the Behavioural Variant of Frontotemporal Dementia. *Brain*, 134(9), 2456–2477.
- Ratnavalli, E., Brayne, C., Dawson, K., & Hodges, J. R. (2002). The Prevalence of Frontotemporal Dementia. *Neurology*, 58(11), 1615–1621.
- Regenbogen, Christina, et al. (2012). Multimodal Human Communication - Targeting Facial Expressions, Speech Content and Prosody. *NeuroImage*, 60(4), 2346–2356. <https://doi.org/10.1016/j.neuroimage.2012.02.043>.
- Reniers, Renate L. E. P., Völlm, Birgit A., Elliott, Rebecca, & Corcoran, Rhiannon (2014). Empathy, ToM, and Self-Other Differentiation: An fMRI Study of Internal States. *Social Neuroscience*.
- Rilling, James K., et al. (2008). Social Cognitive Neural Networks during In-Group and Out-Group Interactions. *NeuroImage*, 41(4), 1447–1461.
- Rogalsky, Corianne, Rong, Feng, Saberi, Kourosh, & Hickok, Gregory (2011). Functional Anatomy of Language and Music Perception: Temporal and Structural Factors Investigated Using Functional Magnetic Resonance Imaging. *The Journal of Neuroscience*, 31(10), 3843–3852. <http://www.jneurosci.org/content/31/10/3843.abstract>.
- Rosen, H. J., et al. (2002). Patterns of Brain Atrophy in Frontotemporal Dementia and Semantic Dementia. *Neurology*, 58(2), 198–208.
- Rosenblau, Gabriela, et al. (2016). The Role of the Amygdala in Naturalistic Mentalising in Typical Development and in Autism Spectrum Disorder. *British Journal of Psychiatry*, 208(6), 556–564.
- Roser, Patrik, et al. (2012). Alterations of Theory of Mind Network Activation in Chronic Cannabis Users. *Schizophrenia Research*, 139(1–3), 19–26. <https://doi.org/10.1016/j.schres.2012.05.020>.
- Rothmayr, Christoph, et al. (2011). Common and Distinct Neural Networks for False-Belief Reasoning and Inhibitory Control. *NeuroImage*, 56(3), 1705–1713.
- Ruckmann, Judith, et al. (2015). How Pain Empathy Depends on Ingroup/Outgroup Decisions: A Functional Magnetic Resonance Imaging Study. *Psychiatry Research - Neuroimaging*, 234(1), 57–65. <https://doi.org/10.1016/j.pscychresns.2015.08.006>.

- Rutherford, MD, ... S Baron-Cohen - Journal of autism and, and Undefined 2002. 2002. "Reading the Mind in the Voice Original." *Journal of Autism and Developmental Disorders*.
- Sachs, Matthew E., Habibi, Assal, Damasio, Antonio, & Kaplan, Jonas T. (2020). Dynamic Intersubject Neural Synchronization Reflects Affective Responses to Sad Music. *NeuroImage (June 2019)*, Article 116512. <https://doi.org/10.1016/j.neuroimage.2019.116512>.
- Sacks, Oliver (2007). *Musicophilia. Tales of Music and the Brain*. London: Vintage.
- Saft, Carsten, et al. (2013). Mentalizing in Preclinical Huntington's Disease: An FMRI Study Using Cartoon Picture Stories. *Brain Imaging and Behavior*, 7(2), 154–162.
- Salimpoor, Valorie N. et al. 2011. Anatomically Distinct Dopamine Release during Anticipation and Experience of Peak Emotion to Music. In *Nature Neuroscience*, 257–264.
- Samson, Andrea C., Zysset, Stefan, & Huber, Oswald (2008). Cognitive Humor Processing: Different Logical Mechanisms in Nonverbal Cartoons - An FMRI Study. *Social Neuroscience*.
- Saur, D., et al. (2008). Ventral and Dorsal Pathways for Language. In *Proceedings of the National Academy of Sciences* (pp. 18035–18040).
- Saxe, R., & Kanwisher, N. (2003). People Thinking about Thinking People: The Role of the Temporo-Parietal Junction in 'Theory of Mind'. *NeuroImage*.
- Saxe, Rebecca, & Powell, Lindsey J. (2006). It's the Thought That Counts: Specific Brain Regions for One Component of Theory of Mind. *Psychological Science*, 17(8), 692–699.
- Schlaffke, Lara, et al. (2015). Shared and Nonshared Neural Networks of Cognitive and Affective Theory-of-Mind: A Neuroimaging Study Using Cartoon Picture Stories. *Human Brain Mapping*, 36(1), 29–39.
- Schmitgen, Mike M., et al. (2016). Stimulus-Dependent Amygdala Involvement in Affective Theory of Mind Generation. *NeuroImage*, 129, 450–459.
- Schmitzhorst, Vincent J. (2005). Separate Cortical Networks Involved in Music Perception Preliminary Functional MRI Evidence for Modularity of Music Processing. *NeuroImage*, 25(2), 444–451.
- Schnell, Knut, Bluschke, Sarah, Konradt, Brigitte, & Walter, Henrik (2011). Functional Relations of Empathy and Mentalizing: An FMRI Study on the Neural Basis of Cognitive Empathy. *NeuroImage*, 54(2), 1743–1754.
- Schroeter, Matthias L., Racza, Karolina, Neumann, Jane, & Yves von Cramon, D. (2008). Neural Networks in Frontotemporal Dementia-A Meta-Analysis. *Neurobiology of Aging*, 29(3), 418–426.
- Schroeter, Matthias L., Racza, Karolina, Neumann, Jane, & Yves von Cramon, D. (2007). Towards a Nosology for Frontotemporal Lobar Degenerations-A Meta-Analysis Involving 267 Subjects. *NeuroImage*, 36(3), 497–510.
- Schulte-Rüther, Martin, et al. (2008). Gender Differences in Brain Networks Supporting Empathy. *NeuroImage*, 42(1), 393–403.
- Schultz, Robert T., et al. (2003). The Role of the Fusiform Face Area in Social Cognition: Implications for the Pathobiology of Autism. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 358(1430), 415–427.
- Schurz, Matthias, et al. (2014). Fractionating Theory of Mind: A Meta-Analysis of Functional Brain Imaging Studies. *Neuroscience and Biobehavioral Reviews*, 42, 9–34.
- Schuwerk, Tobias, et al. (2017). The RTPJ's Overarching Cognitive Function in Networks for Attention and Theory of Mind. *Social Cognitive and Affective Neuroscience*, 12(1), 157–168.
- Seara-Cardoso, Ana, Viding, Essi, Lickley, Rachael A., & Sebastian, Catherine L. (2015). Affective Resonance in Response to Others' Emotional Faces Varies with Affective Ratings and Psychopathic Traits in Amygdala and Anterior Insula. *Social Neuroscience*.
- Seara-Cardoso, Ana, Viding, Essi, Lickley, Rachael A., & Sebastian, Catherine L. (2015). Neural Responses to Others' Pain Vary with Psychopathic Traits in Healthy Adult Males. *Cognitive, Affective and Behavioral Neuroscience*, 15(3), 578–588.
- Seeley, William W., et al. (2008). Frontal Paralimbic Network Atrophy in Very Mild Behavioral Variant Frontotemporal Dementia. *Archives of Neurology*, 65(2), 249–255.
- Seeley, William W., Zhou, Juan, & Kim, Eun Joo (2012). Frontotemporal Dementia: What Can the Behavioral Variant Teach Us about Human Brain Organization? *Neuroscientist*, 18(4), 373–385.
- Seeley, William W., et al. (2009). Neurodegenerative Diseases Target Large-Scale Human Brain Networks. *Neuron*, 62(1), 42–52.
- Seger, Carol A., Stone, Maria, & Keenan, Julian P. (2004). Cortical Activations during Judgments about the Self and an Other Person. *Neuropsychologia*, 42(9), 1168–1177.
- Shen, Dongchao, et al. 2018. Brain Structural and Perfusion Signature of Amyotrophic Lateral Sclerosis with Varying Levels of Cognitive Deficit. *Frontiers in Neurology*.
- Sheng, Feng, et al. (2014). Task Modulations of Racial Bias in Neural Responses to Others' Suffering. *NeuroImage*.
- Shibata, Midori, et al. (2011). Neural Processing Associated with Comprehension of an Indirect Reply during a Scenario Reading Task. *Neuropsychologia*, 49(13), 3542–3550. <https://doi.org/10.1016/j.neuropsychologia.2011.09.006>.
- Shibata, Midori, Toyomura, Akira, Itoh, Hiroaki, & Abe, Jun ichi (2010). Neural Substrates of Irony Comprehension: A Functional MRI Study. *Brain Research*, 1308, 114–123. <https://doi.org/10.1016/j.brainres.2009.10.030>.
- Simon, Daniela, Craig, Kenneth D., Miltner, Wolfgang H. R., & Rainville, Pierre (2006). Brain Responses to Dynamic Facial Expressions of Pain. *Pain*, 126(1–3), 309–318.
- Singer, Tania, et al. (2004). Empathy for Pain Involves the Affective but Not Sensory Components of Pain. *Science*, 303(5661), 1157–1162.
- Sommer, Monika, et al. (2007). Neural Correlates of True and False Belief Reasoning. *NeuroImage*, 35(3), 1378–1384.
- Specht, Karsten, & Wiggleworth, Philip (2018). The Functional and Structural Asymmetries of the Superior Temporal Sulcus. *Scandinavian Journal of Psychology*, 59 (1), 74–82.
- Spiers, Hugo J., & Maguire, Eleanor A. (2006). Spontaneous Mentalizing during an Interactive Real World Task: An fMRI Study. *Neuropsychologia*, 44(10), 1674–1682.
- Spotorino, Nicola, et al. (2012). Neural Evidence That Utterance-Processing Entails Mentalizing: The Case of Irony. *NeuroImage*, 63(1), 25–39. <https://doi.org/10.1016/j.neuroimage.2012.06.046>.
- Sprengelmeyer, R., Rausch, M., Eysel, U. T., & Przuntek, H. (1998). Neural Structures Associated with Recognition of Facial Expressions of Basic Emotions. *Proceedings of the Royal Society B: Biological Sciences*, 265(1409), 1927–1931.
- Spunt, Robert, & Ralph Adolphs, P. (2014). Validating the Why/How Contrast for Functional MRI Studies of Theory of Mind. *NeuroImage*, 99, 301–3011.
- Spunt, Robert P., Ellsworth, Emily, & Adolphs, Ralph (2017). The Neural Basis of Understanding the Expression of the Emotions in Man and Animals. *Social Cognitive and Affective Neuroscience*, 12(1), 95–105.
- Spunt, Robert P., & Lieberman, Matthew D. (2012). An Integrative Model of the Neural Systems Supporting the Comprehension of Observed Emotional Behavior. *NeuroImage*, 59(3), 3050–3059. <https://doi.org/10.1016/j.neuroimage.2011.10.005>.
- Spunt, Robert P., Satpute, Ajay B., & Lieberman, Matthew D. (2011). Identifying the What, Why, and How of an Observed Action: An fMRI Study of Mentalizing and Mechanizing during Action Observation. *Journal of Cognitive Neuroscience*, 23(1), 63–74.
- Sripada, Chandra Sehkar, et al. (2009). Functional Neuroimaging of Mentalizing during the Trust Game in Social Anxiety Disorder. *NeuroReport*, 20(11), 984–989.
- Suchy, Yana, and James A. Holdnack. 2013. "Assessing Social Cognition Using the ACS for WAIS-IV and WMS-IV." In WAIS-IV, WMS-IV, and ACS: Advanced Clinical Interpretation.
- Suzuki, Shinsuke, et al. (2012). Learning to Simulate Others' Decisions. *Neuron*, 74(6), 1125–1137. <https://doi.org/10.1016/j.neuron.2012.04.030>.
- Takahashi, Haruka K., et al. (2015). Brain Networks of Affective Mentalizing Revealed by the Tear Effect: The Integrative Role of the Medial Prefrontal Cortex and Precuneus. *Neuroscience Research*, 101, 32–43.
- Tamm, Sandra, et al. (2017). The Effect of Sleep Restriction on Empathy for Pain: An fMRI Study in Younger and Older Adults. *Scientific Reports*, 7(1), 1–14.
- Tholen, Matthias G., et al. (2020). Functional Magnetic Resonance Imaging (fMRI) Item Analysis of Empathy and Theory of Mind. *Human Brain Mapping*.
- Thye, Melissa D., Murdaugh, Donna L., & Kana, Rajesh K. (2018). Brain Mechanisms Underlying Reading the Mind from Eyes, Voice, and Actions. *Neuroscience*, 374, 172–186. <https://doi.org/10.1016/j.neuroscience.2018.01.045>.
- Todorov, Alexander, Ida Gobbini, M., Evans, Karla K., & Haxby, James V. (2007). Spontaneous Retrieval of Affective Person Knowledge in Face Perception. *Neuropsychologia*, 45(1), 163–173. <https://doi.org/10.1016/j.neuropsychologia.2006.04.018>.
- Toivainen, Petri, et al. (2014). Capturing the Musical Brain with Lasso: Dynamic Decoding of Musical Features from fMRI Data. *NeuroImage*.
- Trost, Wiebke, Ehofer, Thomas, Zentner, Marcel, & Vuilleumier, Patrik (2012). Mapping Aesthetic Musical Emotions in the Brain. *Cerebral Cortex*, 22(12), 2769–2783.
- Turkeltaub, Peter E., et al. (2012). Minimizing Within-Experiment and within-Group Effects in Activation Likelihood Estimation Meta-Analyses. *Human Brain Mapping*, 33 (1), 1–13.
- Uchiyama, Hitoshi, et al. (2006). Neural Substrates of Sarcasm: A Functional Magnetic Resonance Imaging Study. *Brain Research*, 1124(1), 100–110.
- Ushida, Takahiro, et al. (2008). Virtual Needle Pain Stimuli Activates Cortical Representation of Emotions in Normal Volunteers. *Neuroscience Letters*, 439(1), 7–12.
- Vachon-Presseau, E., et al. (2012). Neural Processing of Sensory and Emotional-Communicative Information Associated with the Perception of Vicarious Pain. *NeuroImage*, 63(1), 54–62. <https://doi.org/10.1016/j.neuroimage.2012.06.030>.
- Vanderwal, Tamara, et al. (2008). Self, Mother and Abstract Other: An fMRI Study of Reflective Social Processing. *NeuroImage*, 41(4), 1437–1446.
- Veroude, Kim, et al. (2012). The Effect of Perspective and Content on Brain Activation during Mentalizing in Young Females. *Journal of Clinical and Experimental Neuropsychology*, 34(3), 227–234.
- Vistoli, Damien, Achim, Amélie M., Lavoie, Marie Audrey, & Jackson, Philip L. (2016). Changes in Visual Perspective Influence Brain Activity Patterns during Cognitive Perspective-Taking of Other People's Pain. *Neuropsychologia*, 85, 327–336. <https://doi.org/10.1016/j.neuropsychologia.2016.03.020>.
- Vogelky, K., et al. (2001). Mind Reading: Neural Mechanisms of Theory of Mind and Self-Perspective. *NeuroImage*, 14(1), 170–181.
- Völlm, Birgit A., et al. (2006). Neuronal Correlates of Theory of Mind and Empathy: A Functional Magnetic Resonance Imaging Study in a Nonverbal Task. *NeuroImage*, 29 (1), 90–98.
- Walter, H., et al. (2009). Dysfunction of the Social Brain in Schizophrenia Is Modulated by Intention Type: An fMRI Study. *Social Cognitive and Affective Neuroscience*, 4(2), 166–176.
- Walter, Henrik, et al. (2004). Understanding Intentions in Social Interaction: The Role of the Anterior Paracingulate Cortex. *Journal of Cognitive Neuroscience*, 16(10), 1854–1863.
- Wang, A. Ting, Lee, Susan S., Sigman, Marian, & Dapretto, Mirella (2006). Developmental Changes in the Neural Basis of Interpreting Communicative Intent. *Social cognitive and affective neuroscience*, 1(2), 107–121.
- Wang, Yi, et al. (2015). Dimensional Schizotypy and Social Cognition: An fMRI Imaging Study. *Frontiers in Behavioral Neuroscience*, 9(MAY), 1–9.
- Whitehead, Jocelyne C., & Armony, Jorge L. (2019). Multivariate fMRI Pattern Analysis of Fear Perception across Modalities. *European Journal of Neuroscience*.
- Whitwell, J. L., et al. (2009). Atrophy Patterns in IVS10+16, IVS10+3, N279K, S305N, P301L, and V337M MAPT Mutations. *Neurology*, 73(13), 1058–1065.

- Whitwell, Jennifer L., et al. (2004). Voxel-Based Morphometry in Tau-Positive and Tau-Negative Frontotemporal Lobar Degenerations. *Neurodegenerative Diseases*, 1(4–5), 225–230.
- Whitwell, Jennifer L., et al. (2005). Magnetic Resonance Imaging Signatures of Tissue Pathology in Frontotemporal Dementia. *Archives of Neurology*, 62(9), 1402–1408.
- Whitwell, J. L., et al. (2011). Imaging Signatures of Molecular Pathology in Behavioral Variant Frontotemporal Dementia. *Journal of Molecular Neuroscience*.
- Wicker, Bruno, et al. (2003). Both of Us Disgusted in My Insula: The Common Neural Basis of Seeing and Feeling Disgust. *Neuron*, 40(3), 655–664.
- Wilson, Nikki Anne, et al. (2020). Scene Construction Impairments in Frontotemporal Dementia: Evidence for a Primary Hippocampal Contribution. *Neuropsychologia*, 137.
- Wilson, Stephen M., et al. (2009). The Neural Basis of Surface Dyslexia in Semantic Dementia. *Brain*, 132(1), 71–86.
- Wilson, Stephen M., et al. (2010). Connected Speech Production in Three Variants of Primary Progressive Aphasia. *Brain*, 133(7), 2069–2088.
- Wolf, Ingo, Dziobek, Isabel, & Hecker, Hauke R. (2010). Neural Correlates of Social Cognition in Naturalistic Settings: A Model-Free Analysis Approach. *NeuroImage*, 49 (1), 894–904. <https://doi.org/10.1016/j.neuroimage.2009.08.060>.
- Wong, Stephanie, et al. (2016). Comparison of Prefrontal Atrophy and Episodic Memory Performance in Dysexecutive Alzheimer's Disease and Behavioral-Variant Frontotemporal Dementia. *Journal of Alzheimer's Disease*, 51(3), 889–903.
- Xiao, Hui, Jacobsen, Andre, Chen, Ziqian, & Wang, Yang (2017). Detecting Social-Cognitive Deficits after Traumatic Brain Injury: An ALE Meta-Analysis of fMRI Studies. *Brain Injury*, 31(10), 1331–1339. <https://doi.org/10.1080/02699052.2017.1319576>.
- Yang, Jing, Pan, Pinglei, Song, Wei, & Shang, Hui Fang (2012). Quantitative Meta-Analysis of Gray Matter Abnormalities in Semantic Dementia. *Journal of Alzheimer's Disease*, 31(4), 827–833.
- Young, Liane, Cushman, Fiery, Hauser, Marc, & Saxe, Rebecca (2007). The Neural Basis of the Interaction between Theory of Mind and Moral Judgment. In *Proceedings of the National Academy of Sciences of the United States of America* (pp. 8235–8240).
- Young, Liane, Dodell-Feder, David, & Saxe, Rebecca (2010). What Gets the Attention of the Temporo-Parietal Junction? An fMRI Investigation of Attention and Theory of Mind. *Neuropsychologia*, 48(9), 2658–2664.
- Young, Liane, Scholz, Jonathan, & Saxe, Rebecca (2011). Neural Evidence for 'Intuitive Prosecution': The Use of Mental State Information for Negative Moral Verdicts. *Social Neuroscience*.
- Zahn, Roland, et al. (2005). Mapping of Temporal and Parietal Cortex in Progressive Nonfluent Aphasia and Alzheimer's Disease Using Chemical Shift Imaging, Voxel-Based Morphometry and Positron Emission Tomography. *Psychiatry Research - Neuroimaging*, 140(2), 115–131.
- Zaitchik, Deborah, et al. (2010). Mental State Attribution and the Temporoparietal Junction: An fMRI Study Comparing Belief, Emotion, and Perception. *Neuropsychologia*, 48(9), 2528–2536. <https://doi.org/10.1016/j.neuropsychologia.2010.04.031>.
- Zamboni, G., et al. (2008). Apathy and Disinhibition in Frontotemporal Dementia: Insights into Their Neural Correlates. *Neurology*, 71(10), 736–742.
- Zatorre, R. J., and V. N. Salimpoor. 2013. From Perception to Pleasure: Music and Its Neural Substrates. *Proceedings of the National Academy of Sciences* 110 (Supplement_2): 10430–37. <http://www.pnas.org/cgi/doi/10.1073/pnas.1301228110>.
- Zhang, Heming, et al. (2018). Facial Expression Enhances Emotion Perception Compared to Vocal Prosody: Behavioral and fMRI Studies. *Neuroscience Bulletin*, 34(5), 801–815. <https://doi.org/10.1007/s12264-018-0231-9>.
- Zheng, Li, Zhang, Fangxiao, et al. (2016). Decreased Empathic Responses to the 'Lucky Guy' in Love: The Effect of Intrasexual Competition. *Frontiers in Psychology*, 7(MAY), 1–8.
- Zheng, Li, Wang, Qianfeng, et al. (2016). Perceived Reputation of Others Modulates Empathic Neural Responses. *Experimental Brain Research*, 234(1), 125–132.
- Zhou, Juan, et al. (2010). Divergent Network Connectivity Changes in Behavioural Variant Frontotemporal Dementia and Alzheimer's Disease. *Brain*, 133(5), 1352–1367.
- Zhou, Juan, & Seeley, William W. (2014). Network Dysfunction in Alzheimer's Disease and Frontotemporal Dementia: Implications for Psychiatry. *Biological Psychiatry*, 75 (7), 565–573.
- Ziae, Maryam, et al. (2016). The Impact of Aging on the Neural Networks Involved in Gaze and Emotional Processing. *Neurobiology of Aging*, 48, 182–194. <https://doi.org/10.1016/j.neurobiolaging.2016.08.026>.